



## OPEN ACCESS

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/injuryprev-2015-041784>).

<sup>1</sup>Faculty of Health and Life Sciences, School of Veterinary Science, University of Liverpool, Leahurst Campus, Cheshire, UK

<sup>2</sup>Faculty of Health and Life Sciences, Department of Epidemiology and Population Health, Institute of Infection and Global Health, University of Liverpool, Leahurst Campus, Cheshire, UK

<sup>3</sup>Institute for Risk and Uncertainty, University of Liverpool

## Correspondence to

Dr Carri Westgarth, Department of Epidemiology and Population Health, Institute of Infection and Global Health, Faculty of Health and Life Sciences, University of Liverpool, Leahurst Campus, Chester High Road, Neston, Cheshire CH64 7TE, UK; [carri.westgarth@liverpool.ac.uk](mailto:carri.westgarth@liverpool.ac.uk)

Received 5 August 2015

Revised 9 December 2015

Accepted 11 December 2015

Published Online First

12 January 2016



CrossMark

**To cite:** Fraser-Williams AP, McIntyre KM, Westgarth C. *Inj Prev* 2016;**22**:437–441.

# Are cattle dangerous to walkers? A scoping review

Angharad P Fraser-Williams,<sup>1</sup> K Marie McIntyre,<sup>2</sup> Carri Westgarth<sup>1,2,3</sup>

## ABSTRACT

Anecdotal evidence suggests that people coming into contact with cattle while participating in outdoor pursuits can sustain severe, even fatal injuries. This has negative implications for farmers, cattle and the public. This study outlines findings from a review of published literature, UK media reports and internet guidelines currently available to the UK public for walking near cattle. A total of 54 cattle attacks were reported in the UK media from 1 January 1993 to 31 May 2013; approximately one-quarter resulted in fatality and two-thirds involved dogs. Walking with dogs among cows, particularly with calves present, was a problematic context. Twenty pieces of commonly occurring advice were found within various guidelines. However, there are no definitive approved guidelines, no published studies describing the prevalence of cattle attacks on members of the public and no system in place to document them. Attacks by cattle are underinvestigated and further work should assess their public health impact.

## INTRODUCTION

Annually, there are 3.6 billion tourist visits to the UK countryside; 18% of visitors being walkers.<sup>1 2</sup> In England in 2011–2012, dog walking was the second most frequent countryside activity, accounting for 51.0% of visits.<sup>3</sup> As promotion of physical activity via dog walking is a public health strategy, evaluation of potential risks to public health from these activities is important.<sup>4</sup>

There are approximately 300 000 UK farms<sup>5</sup> and 9.7 million cattle.<sup>6</sup> Public rights of way cross farmland including livestock grazing, so that while walking people come into contact with livestock, particularly cattle, and may have limited knowledge of how to behave around them.

Risk assessments and mitigation measures are increasingly implemented to manage risk, but intensive attempts to govern human/cattle interactions would have negative implications for farmers, cattle and the public. Likely risk-management practices include removal of herds from publically accessed fields during peak seasons or restricting public access, and euthanasia of cattle that have attacked walkers. These could affect animal welfare via husbandry and farming practices, and financially impact farmers. Further, they could reduce public motivation for countryside outdoor activities. Thus, it is important to fully evaluate the scope of this public health problem.

The aim of this project was to assess available information about negative interactions between the public and cattle, to identify risk factors for cattle attacks, and highlight the availability and usefulness of guidance on walking among livestock. The study outlines the findings of: (1) a literature review of published research; (2) a review of media

reports; (3) an internet search of guidelines for countryside walking.

## METHODS

### Search-term development

Search terms for sets of searches were developed by iteratively trialling popular related phrases found in the published literature, media reports and using internet searches, until the most commonly occurring search terms were identified. Using this method, ‘cow’ for example was identified as being more commonly used than ‘cattle’, and sufficient to return all findings also related to ‘cattle’. The searches were conducted by a single author.

### Published literature

Relevant published literature in English was identified using searches of SCOPUS and Web of Knowledge. Formal literature searches of ‘title’, ‘abstract’, ‘keywords’ and ‘topic’ search fields were undertaken using the ‘(cow\* OR bovine) AND (attack OR injury)’ search term, to May 2013 inclusive (figure 1).

### Media reports

Media reports from the UK newspapers describing cattle attacks were identified by searching Lexis Library (1993 to May 2013 inclusive) using the search phrase ‘(cow\* OR bovine) AND (attack OR injury) AND (walker\* OR public)’.

### Public guidelines

Current publically available internet guidelines describing best practice for walking among cattle were identified by searching Google, timespan unrestricted, UK pages only. The prepiloted search terms included: guidelines for walking through fields with cows; walking through fields with cows; cattle in fields crossed by public access; walking a dog through fields with cows; dog walking in cow fields.

## RESULTS


### Published literature

Eight published papers reported outcomes and/or risk factors for cattle attacks (figure 1). Of these, six studies detailed injuries caused by cattle/bulls via hospital admittance.<sup>7–12</sup> Two studies described bovine behaviour such as maternal defensive aggression/vigilance as a fear response to presence of a person/dog or movement to unfamiliar locations.<sup>13 14</sup>

### Injury or fatality records

Kicking, crushing, head butting and trampling were the main causes of hospital-recorded injury, with fractures/contusions the most common consequences. Most injuries were associated with

## PUBLISHED LITERATURE SEARCH



**Figure 1** A schematic diagram of the search phrases used with the SCOPUS and Web of Knowledge (WoK) bibliometric databases, including the number of records returned during a search of published literature relating to outcomes and risk factors of cattle attacks on people undertaking outdoor pursuits.

occupations involving direct cattle contact as opposed to public walkers, for example, dairy and beef farmers, vets and abattoir workers.<sup>7 8</sup> Fractures were the most common injury, kicking the most common mechanism, and feeding the most common situation.<sup>8</sup> Injuries from bulls were more common than from cows.<sup>9</sup> Specific bull-only incidents were also reported.<sup>10 11</sup> US data identified 287 bull cases; 261 were attacks on people, of which 149 were fatal.<sup>11</sup> The injuries related to being charged or trampled, or accidentally stepped on by Holstein dairy and Angus beef dominant breeds.<sup>11</sup>

### Maternal behaviour

A study of maternal defensive aggression as a result of calves, a heritable trait, suggested that humans are perceived as a threat to calves, causing protective behaviour in cows.<sup>13</sup> Extensification of pasture-based systems, decreased routine contact between handlers and non-lactating cows, an increase in

calf contact post partum (to comply with EU ear-tagging regulations) and an increase, by the public, in recreational use of agricultural land were suggested as factors intensifying injury risk, with authors concluding an increasing trend in cattle maternal defensiveness and associated risk to handlers and public safety.<sup>13</sup>


### Fear of dogs

A study of vigilance to measure fear in dairy cattle concluded that dogs were even more threatening than unfamiliar humans, resembling potential predators, and cattle were exceptionally vigilant in novel locations.<sup>14</sup>

### Media

One thousand records were identified, 100 of which were relevant. Of these, 11 were duplicates, leaving 89 records (see online supplementary data).

**Figure 2** The number of cattle attacks on members of the public out walking in livestock areas reported in the media between 1 January 1993 and 31 May 2013, characterised by whether attacks were fatal and involved a dog.



From January 1993 to May 2013, 54 separate cattle attacks were reported involving the public walking in livestock areas (figure 2): attacks peaked at 13 in 2009, of which nine occurred in May and June. Thirteen (24.1%) attacks resulted in walker fatality, with a peak of four in 2009. Reported walker injuries included fractures to arms, ribs, wrist, scapula, clavicle, legs, lacerations, punctured lung, bruising, black eyes, joint dislocation, nerve damage and unconsciousness.

Most (72%) media reports definitively stated the presence/absence of dogs during an attack: 35 (64.8%) involved one dog, and there was no dog in four (7.4%). Reports also described five cattle categories: herd/multiple cattle (n=26, 48.2%); single cow (n=12, 22.2%); cows and calves (n=11, 20.4%); heifer (n=4, 7.4%); and bull (n=1, 1.9%).

Of the attacks, four were reported in more than five newspapers: one described an attack on David Blunkett, the former Home Secretary, who stumbled and broke a rib while releasing his guide-dog's lead; one described an attack on a veterinarian who was trampled and killed while protecting her dogs, who were thought to have sparked the attack; one described an attack on brothers, one of whom died and the other suffered a punctured lung while walking dogs; and one described an attack on a walker who required emergency neurosurgery. She had strayed from an obstructed public footpath to follow a diversion while dog walking; she picked up the dog after it startled cattle.

Media reports suggest factors influencing the likelihood or severity of attack including: the context as well as the behaviour of the people, dogs and cattle (table 1). Descriptive investigation of the data also suggests that female members of the public may be more likely to protect their dogs, while males let them go (see online supplementary data).

Sensational language is used throughout reports, with phrases such as 'Cow attacks: "It looked like they wanted to kill him;"'<sup>15</sup> such language depicts cattle as being dangerous and highly likely to attack at any point, while the walker is portrayed as blameless.

## Guidelines

Twenty-four separate web pages specifying countryside guidelines were identified, including not hanging onto dogs if threatened by cattle, releasing them; and avoiding getting between cows and calves (table 2). The content of the guidelines varied widely, however guidelines on best practice when dog walking stipulated in 'The Countryside Code'<sup>16</sup> were often reproduced. The Health and Safety Executive (HSE) also provides advice for those grazing cattle on public access fields, recognising the main

risk factors: that walkers often have dogs; and when stressed by the weather, illness, unusual disturbance or when maternal instincts are aroused, cattle can become aggressive.<sup>17</sup>

**Table 1** Potential factors identified from media reports of cattle attacks on people out walking in livestock areas (1 January 1993 to 31 May 2013) which may have influenced the likelihood or severity of an attack

Context	Potential factors identified
Circumstances	Location If public footpath was followed Gender of the walker Age of the walker Walking alone or with others Walking with one or more dogs Dog on or off a lead Dog antagonised the cattle Action of the walkers towards the cattle Cattle type: herd/cattle; cows and calves; cow; heifer; bull
Behaviour of the walker (and dog)	Strayed from the right of way Running/walking/cycling Walked between a mother and calf Fell over Dog possibly sparked the attack Dog ran into herd Owner tried to rescue the dog Owner picked up the dog to protect it Owner let go of the dog's lead Owner put dog on a lead to walk around the cattle
Behaviour of the cattle	Attacked Trampled Charged Protecting calf Tossed person into the air Scared of, interested in or ignored the dog Kicked Stamped Knocked down Crushing Butting Gore Surrounded Startled Irrate Running towards walkers Unknown

**Table 2** The most commonly occurring guidelines and advice obtained when searching for information using the 'Google' search engine, which describes best practice when walking through fields of cattle

Guidelines	Number of mentions	Sources
Leave gates and property as you find them	3	Countryside Code Natural England The Ramblers
Follow paths unless wider access is available such as 'Open Access' land	3	Countryside Code Natural England The Ramblers
Do not be afraid of cattle but be mindful they are protective of their young	1	NFU Cymru
Take a walking stick with you	2	NFU Cymru livefortheoutdoors.com
Be bold and walk straight through the cattle if the animals move towards you	1	NFU Cymru
Avoid getting between cows and their calves. Be prepared for cattle to react to your presence, especially if you have a dog with you	1	NFU and Ramblers
Move quickly and quietly, and if possible walk around the herd	1	NFU and Ramblers
Do not put yourself at risk. Find another way round the cattle and rejoin the footpath as soon as possible	1	NFU and Ramblers
Do not panic or run. Most cattle will stop before they reach you. If they follow, just walk on quietly	1	NFU and Ramblers
If you feel threatened, just carry on as normal, do not run, move to the edge of the field and if possible find another way round the field, returning to the original path as soon as is possible	1	go4awalk.com
Try to keep quiet and move away calmly and out of the field as soon as possible. Try not to surprise the cows	1	go4awalk.com
If cows get too close, turn quietly to face them with arms outstretched	1	go4awalk.com
Walk around cattle, rather than through them	1	livefortheoutdoors.com
If jostled by cows, turn to face them, wave your arms to make yourself look as big as possible and shout firmly	1	livefortheoutdoors.com
Apparently bulls can be controlled by twisting the ring in their nose	2	livefortheoutdoors.com walkingworld.com
If you want cattle to go away, wave your arms around and shout at them	1	walkingworld.com
Bulls: give it a reasonably wide berth and walk as quietly round it as you can (try not to run). If you have a close encounter with one you can (in theory) use the ring in its nose to control it, grab the ring and twist it	1	walkingworld.com
Keep dogs under effective control:	5	Countryside Code Natural England NFU and Ramblers
On a lead, or in sight at all times, be aware of what it is doing and be confident it will return to you promptly on command.		go4awalk.com livefortheoutdoors.com
Ensure it does not stray off the path or area where you have a right of access.		
If cattle chase you and your dog, it is safer to let your dog off the lead—do not risk getting hurt by trying to protect it	6	Countryside Code NFU Cymru NFU and Ramblers The Ramblers livefortheoutdoors.com walkingworld.com
Take extra care when walking dogs around livestock (especially young farm animals)	3	The Ramblers
Be especially vigilant if you find yourself with your dog in a field with both cows and calves		go4awalk.com walkingworld.com

Following three cattle-related deaths in 2009, the National Farmers Union provided new signage reminding walkers using public footpaths to keep dogs on leads, but release if chased or threatened by cattle.<sup>18</sup> In October 2005, the Countryside and Rights of Way Act 2000 (CROW) was implemented across England<sup>19</sup> giving the public freedom to roam without staying on paths; restrictions include keeping dogs on leads near livestock and landowners/tenants excluding or restricting access in certain circumstances.

## DISCUSSION

Walking with dogs and cows with calves were key features of this study; both recognised as risk factors in the published literature, newspaper articles and web pages. Often dogs drew cattle's attention, and owners picked up dogs rather than releasing them.

Further, the enormity of the issue of cattle-related injury is unknown, although hospital records describe injuries mainly in animal husbandry and related occupations. This study confirms that walker injury from cattle is a public health risk worth

future attention. There is no official documentation system, and it is likely that attacks are under-reported. Between April 1996 and March 2006, the HSE investigated 46 incidents,<sup>20</sup> whereas the media review reported in this study identified 17 attacks, and at least one cattle attack requiring hospitalisation was not reported in newspapers nor investigated by HSE (Westgarth, personal communication). The lack of public concern about this risk is surprising considering the political and media concern over comparable animal-related injuries such as bites from dogs; however, cattle are not regarded as 'members of the family'.

In conclusion, numerator information from cattle attacks gaining media attention are available but likely underestimate the true extent and no denominator information is available describing the number of countryside dog walkers; both are necessary to calculate the incidence of attacks and aid examination of risk factors. More research is required, in particular considering changes to countryside access and the promotion of physical activity for health. One possibility is the creation of a highly promoted central database, to encourage self-reporting of cattle attacks. Further, a generic approved guidance document in

leaflet and web form would facilitate better management of this problem.

### What is already known on the subject

- ▶ Many members of the public walk or participate in other outdoor pursuits in the UK countryside and they often have dogs with them.
- ▶ There are anecdotal and hospital reports of cattle causing injury or death to walkers.

### What this study adds

- ▶ Injuries from cattle are a significant and under-reported public health risk.
- ▶ Walkers with dogs are at particular risk.
- ▶ Guidelines of how to behave around cattle and avoid injury vary, in particular concerning control of dogs.

**Acknowledgements** CW's father, was hospitalised after being trampled by cattle in 2012. Thanks for the inspiration.

**Contributors** CW and KMM conceived and designed the study. APF-W assisted with study design, performed the searches, data analysis and drafted the manuscript. CW and KMM revised the manuscript. All authors read and approved the final manuscript.

**Funding** This study received no specific funding. CW is supported by a Medical Research Council Population Health Scientist Research Fellowship (grant number G1002402).

**Competing interests** None declared.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** Summary data from the media report searches are available in the supplementary file.

**Open Access** This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

### REFERENCES

- 1 The Ramblers. Stay safe in the countryside this Easter says NFU and Ramblers. 2013. <http://www.ramblers.org.uk/media-centre/press-releases/2013/march/stay-safe-in-the-countryside-this-easter.aspx>
- 2 National Farmer's Union. Walk safely this Easter says the NFU. 2013. <https://www.nfuonline.com/news/press-centre/stay-safe-in-the-countryside-this-easter/>
- 3 Natural England. Monitor of Engagement with the Natural Environment: The national survey on people and the natural environment—Annual Report from the 2011–12 survey (NECR094). 2012. <http://publications.naturalengland.org.uk/publication/1712385>
- 4 Westgarth C, Christley RM, Christian HE. How might we increase physical activity through dog walking?: A comprehensive review of dog walking correlates. *Int J Behav Nutr Phys Act* 2014;11:83.
- 5 Mulvany P. Secondary CAP's impact on productive structures and family-based agriculture in Europe. 2010. [http://www.ukfg.org.uk/UK\\_CAP\\_casestudy.pdf](http://www.ukfg.org.uk/UK_CAP_casestudy.pdf)
- 6 Defra. Farming Statistics—Livestock Populations at 1 December 2012, UK and England. 2013. [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/183227/defra-stats-foodfarm-landuselivestock-farmstats-dec2012-130314.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/183227/defra-stats-foodfarm-landuselivestock-farmstats-dec2012-130314.pdf)
- 7 Watts M, Meisel EM. Cattle associated trauma—a one year prospective study of all injuries. *Injury* 2011;42:1084.
- 8 Murphy CG, McGuire CM, O'Malley N, et al. Cow-related trauma: a 10-year review of injuries admitted to a single institution. *Injury* 2010;41:548.
- 9 Norwood SMCVVL, Fernandez LG, McLarty JW, et al. Mechanisms and patterns of injuries related to large animals. *J Trauma* 2000;48:740–4.
- 10 Dogan KH, Demirci S, Erkol Z, et al. Injuries and deaths occurring as a result of bull attack. *J Agromedicine* 2008;13:191.
- 11 Sheldon KJ, Deboy G, Field WE, et al. Bull-related incidents: Their prevalence and nature. *J Agromedicine* 2009;14:357.
- 12 Busch HM Jr, Cogbill TH, Landerasper J, et al. Blunt bovine and equine trauma. *J Trauma* 1986;26:559.
- 13 Turner SP, Lawrence AB. Relationship between maternal defensive aggression, fear of handling and other maternal care traits in beef cows. *Livest Sci* 2007;106:182.
- 14 Welp T, Rushen J, Kramer DL, et al. Vigilance as a measure of fear in dairy cattle. *Appl Anim Behav Sci* 2004;87:1.
- 15 Tweedie N. Cow attacks: 'It looked like they wanted to kill him'. 2013. <http://www.telegraph.co.uk/earth/countryside/10064225/Cow-attacks-It-looked-like-they-wanted-to-kill-him.html>
- 16 Natural England. The Countryside Code (NE326). 2012. <http://publications.naturalengland.org.uk/publication/987819?category=38017>
- 17 Health and Safety Executive. Cattle and public access in England and Wales Advice for farmers, landowners and other livestock keepers. 2012. <http://www.hse.gov.uk/pubns/ais17ew.pdf>
- 18 Davies J. New signs issued to farmers to warn walkers of cattle dangers. 2009. <http://www.farmersguardian.com/new-signs-issued-to-farmers-to-warn-walkers-of-cattle-dangers/27350.article>
- 19 Natural England. Open Access land. 2013. <http://www.naturalengland.org.uk/ourwork/access/openaccess/default.aspx>
- 20 Health and Safety Executive. Cattle and Public Access in England and Wales. 2011. <http://adlib.eversite.co.uk/adlib/defra/content.aspx?id=000HK277ZX.0AOWR2SKFLQE8I>

### Who declares Napier a safe city

Napier, New Zealand competed against 30 cities to win the 2016 WHO Western Pacific Regional Office Healthy Cities Recognition for Violence and Injury Prevention (Safe Cities). The city 'demonstrated leadership and collaboration in the fields of, for example, crime prevention, road safety and reducing alcohol-related harm'.

*Comment:* I have never been convinced the Safe City idea actually reduced injuries. But there is no doubting their hard work and dedication.

### Reduction in longevity following a traumatic brain injury?

A Swedish study tracked the life trajectories of a large number of people who sustained a traumatic brain injury (TBI) before the age of 25. TBI victims had shorter life spans than their uninjured siblings and were nearly twice as likely to be hospitalised for a psychiatric illness before age 30. An editorial cautioned that it is not easy to draw a causal connection given the time period.

Key: m – male; f - female

Attack number	Number of papers reporting the attack	Attack date	Location	Public footpath	Walkers details	Category of cattle, as described by the media	Actions of the cattle, as described by the media	Presence of a dog and actions of the owner	Result of the attack
1	5	Nov-10	Nottinghamshire, Stanford on Soar	yes	2: m + f, not residents of the area	bull	Charged, attacked. The bull was later euthanised	N/A	m killed, f critically injured
2	13	Jun-09	Derbyshire, Peak District	N/A	2: m 62, son 26 (David Blunkett, former Home Secretary)	cow	charging, trample, attack, irate	Guide dog, black Labrador-cross, female, put on lead to walk around cattle. Owner tried to release dog but stumbled	broken rib, bruising
3	1	Oct-04	Derbyshire	N/A	f	cows	knocked down, kicked, stamped	N/A	punctured lung, broken ribs, severe bruising to face
4	1	Jul-11	Cornwall, St Dennis	N/A	4: f, m, 2 children	40 dairy cows	running towards the family	Jack Russell, male	Escaped by diving into a bush
5	1	Oct-05	not stated	N/A	3: m + 2 children, cycling	cow	gore the man	N/A	broken ribs, punctured lung, 3 days in intensive care
6	3	May-09	Beaminster	N/A	4: m, f 60, 2 friends	40 cows + calves	surrounded, butted, trampled, charged	2 Labradors, on short leads	broken ribs, stitches to mouth
7	2	Jun-12	County Durham, Romaldkirk	yes	2: m 46, f, holidaymakers	cows	attacked, trampled, reasons unknown	not thought to have a dog	m: serious injuries to head, back, arms, chest, abdomen. f: unhurt
8	2	Oct-10	North Yorkshire,	N/A	3: f 50s + m	cows	knocked to the	2 dogs	f: broken arm +

			Redmire		f 48  separate incidents on same day		ground, trampled	yes	fractured skull, m: minor cuts and bruises. f broken ribs + collapsed lung
9	12	Jun-09	Yorkshire Dales, Pennine Way	N/A	f 49, vet	herd	trapped against wall, trampled	2 Spaniels and a Collie possibly sparked the attack. Owner tried to protect dogs	killed
10	4	Jul-09	Cardiff	no public right of way	f 63	herd, cows	trampled	Dog ran into herd. Owner went amongst the herd to put dog back on lead	killed
11	1	Jul-99	Derbyshire, Peak District	N/A	m 66	3 Charolais cows + calves	charged, attacked, protecting their calves	Golden Retriever, on lead, then let off when cows started charging	killed
12	1	Aug-09	Somerset, Burtle	N/A	m 75, farmer	herd, farmers own	scared by fire engine, trampled	no	killed
13	1	Jun-07	Hereford, Clehonger	yes	m, holidaymaker	herd of cows + calves	attacked: cow was standing on man's chest when he was found	dog	dislocated right knee, fractured left wrist, chest injuries
14	1	Jun-06	Pembrokeshire	yes	f	herd	knocked over, trampled	dog	killed
15	9	May-03	Cumbria, Greystoke	strayed from right of way: footpath blocked	f 45	20 Simmental-cross cows + calves	attacked, tossed in air, scared by dog	Jack Russell, on an extendable lead, startled the cows. Owner picked dog up to protect it	emergency neurosurgery for a head injury, fractured arm + ribs, knee injury
16	2	May-09	Bristol, Wapley Wood	yes	m 42	herd: > 20 cows + calves	trampled	Jack Russell-Spaniel cross, m, on lead. Owner let go of lead during attack	broken ribs, severe bruising, black eyes, large lump on head
17	1	Jul-09	Bath	N/A	m	cow	trampled	N/A	spinal injuries

18	3	Jul-99	Lancashire, Whalley	yes	m 41, postman	heifer + calf	charged, protecting her calf	N/A	severe spinal injuries
19	2	Aug-10	East Sussex, Burwash	yes	f 42	herd, 6 cows	trampled	Schnauzer, on lead. Owner let go of lead	broken scapular, displaced clavicle, 3 broken ribs, nerve damage, hoof mark on knee
20	1	Aug-07	East Sussex, South Downs	N/A	m, Inspector	50 beef cattle	butted, stamped	N/A	4 broken ribs, punctured lung
21	2	Jul-10	Gloucestershire, Forest of Dean	N/A	m 55, holidaymaker	cattle	trampled	Airedale Terrier	serious head injuries
22	1	Jun-12	West Yorkshire, Norwood Green	yes	2: m 49 + son 9	30 to 40 cows	N/A	Dog, on lead, escaped	m unconscious but ok, son escaped
23	1	Sep-10	London, Totteridge	yes	10: pensioners	Friesian cow	N/A	N/A	1 f: broken leg, 2 with severe bruising
24	2	Oct-12	Wiltshire, Chippenham	N/A	f 68	herd	trampled	dog: treated for injuries	killed (cardiac arrest)
25	2	Oct-12	Wiltshire, Cherhill White Horse	yes	f 57, nurse	cow	trampled	2: Rottweiler and Rottweiler-German Shepherd cross, off the lead	extensive bruising, fractured ribs + fractured shoulder blade
26	2	Jun-97	Windermere, Dales Way	yes	m, running	herd of cows	attacked	dog	face sewn back together, lungs drained, 7 broken ribs
27	1	Sep-04	Derbyshire, Little Longstone	yes	f	herd: 6 cows + calves	trampled	dog	punctured lung, 2 broken ribs, bruising to face
28	2	Oct-07	Brighton	yes	m 50	50 + calves	interested in dog, trampled	Golden Retriever, on lead. Owner fell and	life threatening, punctured lung,

								let go of lead. Dog ran away, distracting the cows	broken rib severed an artery
29	1	Jun-09	Suffolk, South Elmham	N/A	f 45	herd of cows	trampled	2. Owner tried to bring dogs under control	killed
30	1	Apr-98	Oxfordshire, Bloxham	N/A	f 40s	herd of cows	trampled	Border Terrier. Owner tried to rescue dog	broken collar bone, punctured lung, broke all 24 ribs
31	1	Oct-07	Bedfordshire, Leighton Buzzard	N/A	m 73, retired doctor	cattle	N/A	N/A	killed
32	2	Sep-10	Cambridgeshire, St Neots Common	N/A	m 59	cows	trampled, attacked	dog	serious chest injuries
33	2	Jul-04	Cumbria, Askam-in-Furness	N/A	f 82	cows + new-born calves	attacked to protect calves from dog	Border Collie	bruising, broken ribs
34	2	Jun-08	Cheadle, Kingsley	yes	2: m 60 + friend	cow	N/A	Spaniel	m: head injury, multiple rib fractures, bruising to back and abdomen
35	1	Jun-09	Scotland	N/A	f	cow	charged	N/A	broken ribs
36	1	May-09	Isle of Man, Colby	yes	f 52	cow	startled	dogs	6 broken ribs, punctured lung
37	1	Jul-10	France, Pyrenees	N/A	f 37	cows: several	trampled	N/A	unconscious
38	1	Jul-10	France, Pyrenees	N/A	m 50, climbing	cow	charged	N/A	head injuries, cuts to arms
39	1	Jun-09	Sussex Downs	N/A	m, police inspector	30 cows	crushed	dog	crushed
40	1	Oct-05	Oxford, Godstow Lock	N/A	2: m 72 + m	brown heifer, among a group of 10 cows + 2 calves	N/A	dog, on lead	used a walking stick and branch to fend off the cow
41	2	Aug-05	Oxford, Godstow	N/A	3: m 41 +	French Saler	N/A	N/A	punctured lung,

			Lock		children; m 14, f 13; cycling	heifer, among a group of 10 cows + 2 calves			broken ribs
42	1	Sep-05	Pembrokeshire	N/A	f 55	40 suckler cows + calves	trampling, crushing, butting	dog	found dead, suffered injuries consistent with trampling, crushing and butting
43	1	Oct-12	Shropshire, Oswestry	N/A	m 46	herd	charged, trampled	dog	face, chest and leg injuries
44	1	Aug-03	Scotland, Kyle of Lochalsh	N/A	m	cattle + calves	N/A	no	died
45	1	Apr-05	Derbyshire, Baslow Edge	N/A	m 58	highland cow	N/A	Dog. Owner walked between mother and calf	6 broken ribs
46	1	Oct-03	Wolvercote, Port Meadow	N/A	f 60	cow	N/A	N/A	2 black eyes, bruising
47	1	Jun-10	Derbyshire, Middleton - by - Youlgreave	N/A	m 78	herd	N/A	N/A	died
48	1	May-09	East Sussex, Guestling	yes	m 34	cows	N/A	Bulldog; catapulted into the air by a cow, died as a result.	broken ribs, bruising
49	1	Nov-99	Dartmoor	N/A	f 31	Black Galloway cow	N/A	2 miniature Dachshunds, on leads tied to owners waist, ignored by cow	leg, head and internal injuries
50	1	Aug-10	East Sussex, Colemans Hatch	N/A	f	Charolais	N/A	no	ankle injury
51	1	Sep-10	Yorkshire, Dales Way	yes	2: m 66 + f	heifer	ignored the dog	3 dogs, let off the lead	bruising, cut to face, suspected cracked rib
52	1	Oct-09	Yorkshire, Dales Way	N/A	f	cow	N/A	N/A	N/A

53	1	May-07	Devon, Culmstock	yes	f 62	cows	trampled	dog	serious chest injuries
54	10	May-13	Wiltshire, Elbow Field	N/A	2: m 65 + m 72	24 adult crossbreed cattle: mixture of Friesian and Limousin	trampled	dogs	m 65: died m 72: collapsed lung, 'stable' condition