What can we learn from international comparisons of social inequalities in road traffic injury mortality?

T H Lu, T L Chiang, J W Lynch

Some contextual effects can be discovered only through international comparisons

In contrast with the plethora of studies concerning the social inequalities in health that have been conducted since the 1980s, there are relatively few related to injuries.^{1,2} There are even fewer studies comparing the influence of these inequalities in different countries. So, the publication of Borrell *et al*'s study on social inequalities in transportation injury mortality across European countries³ is most welcome and timely [see page 138]. Their study elucidated the effect of differing national contexts on the influence of social disparities on road traffic injury (RTI) mortality.

In this commentary, we will first explain what a so-called "contextual effect" is and then illustrate that many such effects can be discovered only through international comparisons. We contend that "dissimilarities" may provide more useful information than "similarities". Because the contextual determinants of social inequality in RTI mortality *between* and *within* countries are different, they have different implications for injury prevention.

WHAT IS A CONTEXTUAL EFFECT?

In linguistics, context refers to the text surrounding a word, giving a better understanding of what the word means. In art, contextualism refers to the way a work of art may only be understood by knowing the historical, political, or cultural circumstances when it was produced. In health research, context similarly refers to the wider situation surrounding the association between an exposure and outcome and how this wider situation may confer meaning on that association.⁴

The study of the effects of collective or group characteristics on individual level outcomes is termed contextual or multi-level analysis. The effects of group level properties on individual level outcomes are contextual effects.⁵ Incorporating group level variables in contextual analysis is important because this approach provides information not captured by individual level data.⁶ For example,

national income level gives information different from individual level income. The national income level may be a marker for countrywide factors potentially related to RTI, such as road conditions, user patterns, or medical care. Such factors affect everyone in the country regardless of their income. Similarly, the national unemployment level may be a marker of the effects of globalization, which affects everyone in a country, employed or unemployed.

Macro level variables, such as political regime, urban planning, road design, regulations, population density, traffic volume, and so on, affect individuals directly and also constrain the choices they make.⁶ Only through international comparisons can these contextual RTI risk factors in any one country be revealed. This kind of information could not be derived from individual level studies conducted within a country. If everyone in a country was exposed to the same contextual risk factors, there would be no interpersonal variation among study subjects.⁷⁻⁹

"SIMILARITIES" OR "DISSIMILARITIES"?

In their international comparisons, Borrell *et al* concentrated mostly on "similarities"—for example, in all countries less educated men had higher death rates compared with better educated men. More might have been learned had they examined why the magnitude of social inequalities differed in different countries. However, a better understanding of "dissimilarities" between countries could provide more information for injury prevention. Each country has its own historical and cultural context that shapes social inequalities in health and this is why Kunitz argued for "particularism" to understand the social determinants of population health.¹⁰

For example, a recent study revealed that in China, higher socioeconomic position (SEP) was related to a less healthy lifestyle, whereas in the United

States, higher SEP was related to a healthier lifestyle.¹¹ The contrast was partially due to cultural factors and partially to level of economic development.

Using Borrell *et al*'s data, we found the magnitude of relative inequality in RTI mortality between countries to be larger than the relative social inequality within a country.³ The age adjusted death rate ratios (RR) among men with low, middle, and high educational achievement ranged from 1.09 in Spain (Madrid) to 1.66 in Switzerland. However, the RR between Madrid and Finland was about 3.0, regardless of educational level. Thus, in explaining inequalities in RTI mortality rates, social contextual differences between countries are more important than social group differences in any one country.

We have simplified and modified Borrell *et al*'s figure 1 to illustrate these differences. The figure shows that the slope of social inequality between countries was steeper than the slope within countries. Though country C has the lowest RTI mortality compared with country A or B, country C has the largest social inequality compared with A or B. Why?

Determinants of social inequality in RTI mortality between countries

To explain these findings, we used Macintyre's conceptualisation.¹² She suggested three explanations for geographical variations in health: (1) compositional explanations, which draw attention to the characteristics of individuals concentrated in particular places; (2) contextual explanations, which focus on structures in the local physical and social environment; (3) collective explanations, which concentrate on sociocultural and historical features of communities.

Macintyre gave an example relevant to injury prevention. Children in deprived areas may not play outdoors because their families do not have gardens or the resources to take them to parks (a compositional resource based explanation). Or they may not do so because too few parks are provided or there are no easy public transport links to parks that do exist (a contextual resource based explanation). She also suggests that within the prevailing culture either play is not seen as important to children or it is not considered safe for children to play with other (strange) children in public places (a collective explanation).

Analogously, high RTI mortality in a country might be because of a higher percentage of dangerous drivers (a compositional explanation); poorer road infrastructure or post-crash care (a

Figure 1 Hypothetical example to illustrate social inequality in road traffic injury mortality *between* and *within* countries (SEP, socioeconomic position).

contextual explanation); or lower standards of safety—that is, less thorough policy implementation and enforcement (a collective explanation).

Determinants of social inequality *within* countries

To explain social inequality in RTI within countries, Laflamme proposed a model that illustrates upstream (social context) and downstream (individual level) mechanisms.^{1–13} At an individual level, the model depicts a pathway from social position, through exposure, to health hazards/risk factors, and on to health outcome. Two downstream mechanisms—differential exposure and differential susceptibility—may come into play in the relation between social position and health outcome.

Contextual influences are highlighted by four entry points, each of which refers to different mechanisms on social stratification, on differential exposure, on different susceptibility, and directly on health.

Finally, Lynch and Kaplan propose a Marxian-Weberian view of how socioeconomic position affects health.¹⁴ Some examples pertaining to RTI are seen in table 1 (see <http://www.injuryprevention.com/supplemental> for table 1). This framework reminds us not to overlook the unequal power relationships that affect the implementation or effectiveness of traffic safety policies. We combined both the above frameworks in table 2 to contrast the determinants of social inequality *between* and *within* countries.

CONCLUSIONS

International comparisons provide a prism through which the wider variations in inequalities between countries come into focus. These comparisons highlight the importance of contextual effects on social inequalities in RTI. Individual risky behaviors are embodied in the social context and the unequal power relationships that affect the adoption of countermeasures, as well as acting to shape different exposures and resources experienced across social groups. Clarifying the determinants of social inequality in RTI mortality *between* and *within* countries will help us develop better injury prevention strategies.

Injury Prevention 2005;11:131–133.
doi: 10.1136/ip.2005.008631

Table 1 can be viewed on our website.

Authors' affiliations

T H Lu, T L Chiang, Institute of Health Policy and Management, School of Public Health, National Taiwan University, Taipei, Taiwan
J W Lynch, Center for Social Epidemiology and Population Health, University of Michigan, USA

Correspondence to: Dr T-H Lu, Institute of Public Health, College of Medicine, National Cheng Kung University, No 1 Dah Hsueh Road, Tainan 701, Taiwan; robertlu@mail.ncku.edu.tw

REFERENCES

- 1 Laflamme L, Diderichsen F. Social differences in traffic injury risks in childhood and youth—a literature review and a research agenda. *Inj Prev* 2000;6:293–8.
- 2 Cubbin C, Smith GS. Socioeconomic inequalities in injury: critical issues in design and analysis. *Annu Rev Public Health* 2002;23:349–75.

Table 2 Scenarios explaining social inequality in road traffic injury (RTI) mortality *between* and *within* countries

	Determinants of social inequality in RTI mortality <i>between</i> countries	Determinants of social inequality in RTI mortality <i>within</i> countries
National social context	<i>Scenario 1: host and vector</i> In face of global economic recession, country A had higher unemployment rate than country C	Less educated and manual workers were more likely to be fired during economic recession in country C
RTI related context	Unemployed families could not afford to buy protective devices, better cars, and health insurance	Unemployed families could not afford to buy protective devices, better cars and health insurance
Individual exposure and resources	More people in country A were vulnerable to severe injury in traffic crashes than country C	People with low socioeconomic position were more vulnerable to severe injury in traffic crashes
National social context	<i>Scenario 2: physical environment</i> Country C had higher income than country A	Country C had larger income inequality than country A
RTI related context	Country C invested more on better road infrastructure than country A	Distribution of resources in maintaining better road infrastructure in country C was unequal
Individual exposure and resources	All people living in country C had exposure to safer road conditions than people live in country A	People living in deprived areas of country C had exposure to more unsafe road infrastructure
National social context	<i>Scenario 3: social environment</i> Country C had higher safety standards than country A	The quality of governance in country C was unequal
RTI related context	Country C adopted speed limits and alcohol impairment laws, but not in country A	Speed limits and alcohol impairment laws were more strictly enforced in better-off areas
Individual exposure and resources	All people living in country C have less exposure risk than people living in country A	People living in deprived areas have higher exposure risk than their counterparts in better-off areas in country C

- 3 **Borrell C**, Plasencia A, Huisman M, *et al*. Education level inequalities and transportation injury mortality in the middle aged and elderly in European settings. *Inj Prev* 2005;**11**:138–42.
- 4 **McLaren L**, Hawe P. Ecological perspectives in health research. *J Epidemiol Community Health* 2005;**59**:6–14.
- 5 **Diez-Roux AV**. A glossary for multilevel analysis. *J Epidemiol Community Health* 2002;**56**:588–94.
- 6 **Diez-Roux AV**. Bringing context back into epidemiology: variables and fallacies in multilevel analysis. *Am J Public Health* 1998;**88**:216–22.
- 7 **Schwartz S**, Carpenter KM. The right answer for the wrong question: consequences of type III error for public health research. *Am J Public Health* 1999;**89**:1175–80.
- 8 **Lu TH**. International comparisons do help and are essential for avoiding type III error in injury prevention research. *Inj Prev* 2001;**7**:270–1.
- 9 **Lu TH**. Earthquake and suicide: bringing context back into disaster epidemiological studies. *Int J Epidemiol* 2004;**33**:1406–9.
- 10 **Kunitz SJ**. The value of particularism in the study of the cultural, social, and behavioral determinants of mortality. In: Chen LC, Kleinman A, Ware NC, eds. *Health and social change in international perspective*. Boston, MA: Harvard University Press, 1994:225–50.
- 11 **Kim S**, Symons M, Popkin BM. Contrasting socioeconomic profiles related to healthier lifestyles in China and the United States. *Am J Epidemiol* 2004;**159**:184–91.
- 12 **Macintyre S**, Ellaway A, Cummins S. Place effects on health: how can we conceptualise, operationalise and measure them. *Soc Sci Med* 2002;**55**:125–39.
- 13 **Lafamme L**. Explaining socio-economic differences in injury risks. *Inj Control Saf Promt* 2001;**8**:149–53.
- 14 **Lynch J**, Kaplan GA. Socioeconomic position. In: Berkman L, Kawachi I, eds. *Social epidemiology*. Oxford: Oxford University Press, 2000:13–35.

Clinical Evidence—Call for contributors

Clinical Evidence is a regularly updated evidence-based journal available worldwide both as a paper version and on the internet. *Clinical Evidence* needs to recruit a number of new contributors. Contributors are healthcare professionals or epidemiologists with experience in evidence-based medicine and the ability to write in a concise and structured way.

Areas for which we are currently seeking authors:

- Child health: nocturnal enuresis
- Eye disorders: bacterial conjunctivitis
- Male health: prostate cancer (metastatic)
- Women's health: pre-menstrual syndrome; pyelonephritis in non-pregnant women

However, we are always looking for others, so do not let this list discourage you.

Being a contributor involves:

- Selecting from a validated, screened search (performed by in-house Information Specialists) epidemiologically sound studies for inclusion.
- Documenting your decisions about which studies to include on an inclusion and exclusion form, which we keep on file.
- Writing the text to a highly structured template (about 1500–3000 words), using evidence from the final studies chosen, within 8–10 weeks of receiving the literature search.
- Working with *Clinical Evidence* editors to ensure that the final text meets epidemiological and style standards.
- Updating the text every six months using any new, sound evidence that becomes available. The *Clinical Evidence* in-house team will conduct the searches for contributors; your task is simply to filter out high quality studies and incorporate them in the existing text.
- To expand the topic to include a new question about once every 12–18 months.

If you would like to become a contributor for *Clinical Evidence* or require more information about what this involves please send your contact details and a copy of your CV, clearly stating the clinical area you are interested in, to Klara Brunnhuber (kbrunnhuber@bmjgroup.com).

Call for peer reviewers

Clinical Evidence also needs to recruit a number of new peer reviewers specifically with an interest in the clinical areas stated above, and also others related to general practice. Peer reviewers are healthcare professionals or epidemiologists with experience in evidence-based medicine. As a peer reviewer you would be asked for your views on the clinical relevance, validity, and accessibility of specific topics within the journal, and their usefulness to the intended audience (international generalists and healthcare professionals, possibly with limited statistical knowledge). Topics are usually 1500–3000 words in length and we would ask you to review between 2–5 topics per year. The peer review process takes place throughout the year, and our turnaround time for each review is ideally 10–14 days.

If you are interested in becoming a peer reviewer for *Clinical Evidence*, please complete the peer review questionnaire at www.clinicalevidence.com or contact Klara Brunnhuber (kbrunnhuber@bmjgroup.com).