0327 **COMPARISON OF MORTALITY FOLLOWING HOSPITALISATION FOR ISOLATED HEAD INJURY IN THE UK AND VICTORIA, AUSTRALIA**

B J Gabbe*, R A Lyons, F E Lecky, O Bouamra, M Woodford, T J Coats, P A Cameron Correspondence: School of Public Health and Preventive Medicine, Monash University, The Alfred Hospital Commercial Rd, Melbourne, Victoria, 3004, Australia

10.1136/ip.2010.029215.327

Background Head injury remains a leading cause of death and disability worldwide but advances in trauma systems, pre-hospital care, and critical care have led to improvements in survival.

Aims Compare isolated head injury mortality in England/Wales with Victoria, Australia.

Methods Population-based, isolated head injury admissions data were available for July 2005 to June 2006. To enable better risk-adjustment, severe isolated head injury (AIS severity >3) data were obtained for the same period from the trauma registries for England/Wales (TARN) and Victoria (VSTR).

Results In 2005–2006, there were 7155 isolated head injury admissions in England/Wales and 1217 in Victoria. The in-hospital mortality was 12% for England/Wales and 9% for Victoria. Adjusting for key confounders, the odds of death in England/Wales were higher (AOR 2.0, 95 CI 1.6 to 2.5). 699 severe isolated head injuries were captured by the VSTR, and 575 by TARN over the 12-month period. The in-hospital mortality was 23% for TARN and 17% for VSTR. After adjusting for key confounders, severely isolated head injury patients were more likely to die in England/Wales than Victoria (AOR 1.6, 95% CI 1.1 to 2.3). 72% of TARN cases were managed at a neurosurgical centre compared to 87% of VSTR cases.

Conclusion Death following hospitalisation for an isolated head injury was higher for England/Wales when compared to Victoria, Australia. Differences in the rate of care at neurosurgical centres appeared to be a key explanatory factor.