had medial collateral ligament injury; three (1.5%) had cruciate ligament injury, and two (1.0%) had lateral collateral ligament injury). We also often see patients with knee injury resulting from tai chi exercise in our clinical practice.

We therefore believe that there is a high potential risk of knee injury during tai chi exercise. While performing such exercise, people should pay attention to protecting the knee joint, and older people with osteoporosis should not perform tai chi at all. Appropriate training intensity should be maintained, and the correct training methods should be used. When carrying out tai chi exercises, it is particularly important to avoid repeated use of the squat to protect the knee joint.

Hong-lin Chen,¹ Kun Liu,² Qin-Sheng You²

¹Nantong University, Nantong, Jiangsu, China;

²Department of Cardiothoracic Surgery, Affiliated Hospital of Nantong University, Nantong, Jiangsu, China

Correspondence to Hong-lin Chen, School of Nursing, University of Nantong, Nantong, Jiangsu Province, PR China; pphss@126.com

Competing interests None.

Provenance and peer review Not commissioned; internally peer reviewed.

Published Online First 11 April 2011

Injury Prevention 2011;**17**:286–287.
doi:10.1136/injuryprev-2011-040027

REFERENCES

1. Day L, Finch CF, Harrison JE, *et al.* Modelling the population-level impact of tai-chi on falls and fall-related injury among community-dwelling older people. *Inj Prev* 2010;**16**:321–6.
2. Wu G, Millon D. Joint kinetics during Tai Chi gait and normal walking gait in young and elderly Tai Chi Chuan practitioners. *Clin Biomech* 2008;**23**:787–95.
3. Zhao SY, Zhang XD, Qi WH, *et al.* Reasons for knee joint pain caused by Yang's shadowboxing. *J Chin Clin Rehabil* 2005;**9**:182–3. Chinese.
4. Cheng CF, Pan XG. Knee injury after tai chi exercise on college students. *J Pet Educ Inst Xing* 2006;**9**:103–5. Chinese.

CORRECTION

doi:10.1136/ip.2008.020495corr1

Mutto M, Lett R, Lawoko S, *et al.*
Intentional Injuries among Ugandan

Youth: a trauma registry analysis. *Inj Prev* 2010;**16**:333–6. doi:10.1136/ip.2008.020495.

- The 2002 spatial distribution of Uganda's population as appears in the October 2010 issue (at the end of the Settings part in the Methods section on page 2 of 4) is incorrect. The correct proportions should be: **32.2%** in central, **30.2%** in eastern, **15.6%** in northern and **22.0%** in western regions.
 - Absolute prevalence difference between males and females should be **34.6%** and not 70.6% as appears in the publication (line 7 of the Discussion section).
 - Intentional injuries among youth constitute **7.2%** of total injury burden in the five regions, not one fifth of injury burden as indicated in the Discussion section.
- The journal apologises for these errors.

Calendar

Conferences and events of interest

doi:10.1136/injuryprev-2011-040108

Edited by Barry Pless

3rd European Injury Prevention Conference

16–17 June 2011, Budapest.

20th International Safe Communities Conference

6–9 September 2011, Falun, Sweden.

19th World Congress on Safety and Health at Work

11–15 September 2011, Istanbul, Turkey.

55th Annual Conference of Association for the Advancement of Automotive Medicine (AAAM)

2–5 October 2011, Paris, France.

5th National Occupational Injury Research Symposium (NOIRS)

18–20 October 2011, Morgantown, West Virginia, USA.

Canadian Conference on Injury Prevention and Safety Promotion

16–18 November 2011, Vancouver, Canada.

Australasian Road Safety Research, Policing and Education Conference 2011

6–9 November 2011, Perth, WA, Australia.

11th World Conference on Injury Prevention and Safety Promotion

1–5 October 2012, Wellington, New Zealand.