

Changes in traffic crash mortality rates attributed to use of alcohol, or lack of a seat belt, air bag, motorcycle helmet, or bicycle helmet, United States, 1982–2001

P Cummings, F P Rivara, C M Olson, K M Smith

Injury Prevention 2006;12:148–154. doi: 10.1136/ip.2005.010975



Four appendices to this article are available on our website.

See end of article for authors' affiliations

Correspondence to:
Dr P Cummings, 250
Grandview Drive, Bishop
CA 93514, USA; peterc@
u.washington.edu

Accepted
17 December 2005

Objective: To estimate the contributions of five risk factors to changes in US traffic crash mortality: (1) alcohol use by drivers and pedestrians, (2) not wearing a seat belt, (3) lack of an air bag, (4) not wearing a motorcycle helmet, and (5) not wearing a bicycle helmet.

Design: Longitudinal study of deaths; attributable deaths were estimated using data from other studies.

Setting: US traffic crashes in 1982–2001.

Subjects: People who died in a crash.

Main outcome measures: Counts of deaths attributed to each risk factor, change in rates of deaths, and counts of lives saved by changes in risk factor prevalence.

Results: There were 858 741 traffic deaths during the 20 year period. Estimated deaths attributed to each factor were: (1) alcohol use, 366 606; (2) not wearing a seat belt, 259 239; (3) lack of an air bag, 31 377; (4) no motorcycle helmet, 12 095; (5) no bicycle helmet, 10 552. Over the 20 years, mortality rates attributed to each risk factor declined: alcohol by 53%; not wearing a seat belt by 49%; lack of an air bag by 17%; no motorcycle helmet by 74%; no bicycle helmet by 39%. There were 153 168 lives saved by decreased drinking and driving, 129 297 by increased use of seat belts, 4305 by increased air bag prevalence, 6475 by increased use of motorcycle helmets, and 239 by increased use of bicycle helmets.

Conclusions: Decreased alcohol use and increased use of seat belts were associated with substantial reductions in crash mortality from 1982 through 2001. Increased presence of air bags, motorcycle helmets, and bicycle helmets were associated with smaller reductions.

Traffic crash mortality rates in the United States declined from 26.4 per 100 000 person-years in 1969 to 14.7 in 2003.^{1,2} At least five factors may have influenced this decline in mortality. Citizen groups that influenced attitudes and laws regarding drunk driving have been credited with part of this decrease.^{3–6} The first state seat belt law was adopted in 1984;^{6–8} by 2004, 80% of 52 000 observed front seat motorists were belted.⁹ By 1994 most new cars had dual air bags and these were required in cars by 1998 and in light trucks by 1999.¹⁰ Observed helmet use by motorcyclists was 63% in 1994, rose to 71% in 2000, but declined to 58% by 2002.¹¹ Bicycle helmet use has become more common in some regions.¹²

We studied US traffic deaths during 1982–2001 to assess the contribution of alcohol use, not wearing a seat belt, lack of an air bag, not wearing a motorcycle helmet, and not wearing a bicycle helmet. We estimated the count and proportion of deaths attributed to each risk factor, the change in prevalence for each factor, and trends in mortality rates for all US traffic crash deaths and for deaths attributed to each of the five risk factors and combinations of these factors.

SUBJECTS AND METHODS

Crash data

Since 1975, the National Highway Traffic Safety Administration has maintained the Fatality Analysis Reporting System (FARS) which has data from all crashes on US public roads with a death within 30 days.¹ These are public data and we were exempted from human subjects review.

Study subjects

Subjects were persons in vehicles (cars, trucks of all kinds, minivans, buses, and all others aside from motorcycles),

riders on motorcycles, and bicyclists and pedestrians who died in a crash with a motor vehicle. Drivers could be in vehicles or on motorcycles.

Risk factors

Alcohol use by each vehicle driver, motorcycle driver, pedestrian, or bicyclist could influence their risk of a deadly crash. Driver alcohol use could also influence the risk of death for other people in the same crash: passengers in the same vehicle as the driver, people in other vehicles, motorcyclists, pedestrians, and bicyclists. Not wearing a seat belt was considered to be a risk factor only for those who died in a vehicle; motorcyclists, pedestrians, and bicyclists cannot wear seat belts. Lack of an air bag was treated as a risk factor only for front seat occupants of vehicles that customarily have air bags—for example, bus drivers were not considered to be at risk for lack of an air bag. Failure to wear a helmet was treated as a risk factor only for motorcyclists and bicyclists.

Attributable fractions and attributed deaths

We estimated the fraction of each death that could be attributed to each risk factor: the attributable fraction [AF] = (risk ratio – 1)/(risk ratio).¹³ For occupants of the same vehicle or motorcycle, we assigned each the AF for the driver's blood alcohol level. When more than one vehicle was involved, all occupants were assigned the two largest AF values for involved drivers: combined AF = 1 – [(1 – AF1) × (1 – AF2)]. For pedestrians and bicyclists we combined their AF and the largest AF of any involved driver. For other

Abbreviation: FARS, Fatality Analysis Reporting System.

Table 1 Traffic crash deaths attributed to each risk factor, US, 1982–2001

Risk factor	Affected population	All deaths in affected population	Deaths attributed to each risk factor	
		Count	Count	Rate* (%)†
Alcohol	All traffic deaths	858 741	366 606	7.15 43
No seat belt	Vehicle occupants	641 114	259 239	5.06 40
No air bag	Vehicle front seat occupants	564 124	31 377	0.61 6
No motorcycle helmet	Motorcycle riders	63 811	12 095	0.24 19
No bicycle helmet	Bicycle riders	16 473	10 552	0.21 64
All five risk factors	All traffic deaths	858 741	528 105‡	10.30 61

*Rates per 100 000 person-years.
 †(%) = 100 × attributed deaths/deaths in affected population.
 ‡Count attributed to all five risk factors is less than the sum of the five individual counts because many persons were at risk from more than one factor.

risk factors, such as not wearing a seat belt, we used the AF based upon the risk factor status of each dead individual.

The risk ratios for alcohol use (see <http://www.injury-prevention.com/supplemental> for Appendix 1) were from a case control study (table 6.6 of that report) that used FARS and other data to estimate the association of blood alcohol level with driver involvement in a passenger vehicle crash.¹⁴ We assumed that these risk ratios could be applied to motorcycle drivers, pedestrians, and bicyclists. The risk ratios for air bags and seat belts came from a matched cohort study that used FARS data; the risk ratios for these two factors were adjusted for each other and an interaction term between air bags and seat belts was used.¹⁵ The estimates for motorcycle helmet use were from a matched cohort study of FARS data.¹⁶ The risk ratios for bicycle helmets were from a case control study.¹⁷ For estimates that involved more than one of these risk factors, we used the formula described above to combine attributable fractions.

For example, consider a dead, unbelted, 32 year old male car driver with an air bag and a blood alcohol of 0.06. A 27 year old female driver in the same crash had a blood alcohol of 0.08. Using Appendix 1, the risk ratio for the dead driver’s alcohol level was 3.76, the risk ratio for the other driver’s alcohol level was 6.25, and the risk ratio for being unbelted with an air bag was 2.33. Lack of a helmet made no contribution to the death. Therefore, the fraction of this driver’s death attributable to all five risk factors was 1

$$- [(1 - (3.76-1)/3.76) \times (1 - (6.25-1)/6.25) \times (1 - (2.33-1)/2.33)] = 0.98.$$

For each year the count of deaths attributed to each risk factor was obtained by summing the relevant attributable fractions. Other deaths not attributed to each risk factor were estimated by subtracting attributed deaths from all deaths.

Changes in risk factor prevalence

We used linear regression to estimate the mean change over time in risk factor prevalence among the dead in the FARS data.¹⁸ Change estimates for air bags were limited to the years 1992–2001, as air bags were rare before 1992. For bicycle helmets we limited change estimates to 1994–2001, as bicycle helmet use was not recorded before 1994 in FARS data; we assumed that no bicycle helmets were used before 1994, as only 2% of dead bicyclists were helmeted in 1994.

Rates, rate changes, and lives saved

To estimate mortality rates (per 100 000 person-years) we used FARS counts of deaths and population estimates from the US Census Bureau.¹⁹ We used Poisson regression, with person-years as an offset, to estimate the change in rates over the 20 years.^{20 21} We estimated rate changes for: (1) deaths attributed to each risk factor, (2) other deaths not attributed to each risk factor, and (3) all deaths, which can be thought of as a weighted average of changes (1) and (2).




Figure 1 Changes in the prevalence of five risk factors among persons who died in traffic crashes in the US during 1982–2001. Alcohol was defined for this figure as any level of blood alcohol by a driver, pedestrian, or bicyclist in the crash. Trends in proportions are shown with fitted lines.

Table 2 Change in risk factor prevalence: percent of the dead exposed to each of five risk factors.* Estimates for 1982 and 2001 are from a line fitted to the data for all years. †

Risk factor	Prevalence in 1982 (%)	Prevalence in 2001 (%)	Difference (%)	95% CI for difference (%)
Alcohol	56	36	-20	-20 to -19
No seat belt	94	55	-39	-40 to -39
No air bag	100	53	-48	-48 to -47
No motorcycle helmet	60	44	-16	-18 to -15
No bicycle helmet‡	100	94	-4	-6 to -3

*Only some dead people were considered to be at risk for some factors: for example, only vehicle occupants (not motorcyclists) were considered at risk for not wearing a seat belt and only motorcycle riders were considered at risk for not wearing a motorcycle helmet.

†Change difference = final year prevalence minus first year prevalence. Change estimate restricted to 1992–2001 for air bags and 1994–2001 for bicycle helmets.

‡Failure to wear a bicycle helmet assumed to be 100% before 1994.

For each risk factor, we predicted the count of deaths that would have occurred in each year if the rate change for all deaths had been the same as the change for deaths not attributed to that factor. In each year, the predicted count of deaths minus the observed count of deaths was an estimate of lives saved (if positive) or lost (if negative) by changes in risk factor prevalence over time. We added these count differences over the 20 years, to estimate lives saved or lost due to changes in risk factor prevalence. These estimates were limited to 1992–2001 for air bags and 1994–2001 for bicycle helmets.

Missing data

Data for blood alcohol concentration were missing for 57% of drivers and 48% of pedestrians and bicyclists. Among those who died, seat belt use was missing for 11% of vehicle occupants, helmet use for 10% of motorcyclists and 0.5% of bicyclists, and age for 0.4%.

Information about air bag presence was in FARS data, but it was often missing. We used software to link the partial vehicle identification number to manufacturer supplied information about air bags.²² We assumed the manufacturer's air bag information was correct, but when it was missing we used information about air bag presence if it was in FARS. After making some assumptions about air bag presence (for example, air bags were not present in buses), air bag presence was missing for 0.2% of vehicle occupants who died.

We used multiple imputation to create 10 sets of data which were identical in regard to known information, but could differ, one from another, on imputed values for missing information.^{23–28} Details of the imputation process can be found elsewhere.^{29–31}

Confidence intervals for changes

Confidence intervals (CI) were from linear or Poisson regression with a robust (sandwich) variance estimator.^{32–33} We generated estimates in each imputed data set, combined the 10 estimates by averaging on the appropriate scale, and computed confidence intervals that accounted for the variance within each data set and the variance between the 10 data sets.³⁴

RESULTS

Deaths attributed to the five study risk factors

There were 858 741 traffic deaths (rate 16.75) during the 20 year period. When the five risk factors were considered separately, 43% (366 606) of all crash deaths could be attributed to alcohol (table 1), 30% (259 239) to not wearing a seat belt, 4% (31 377) to lack of an air bag, and 1% each to lack of a motorcycle (12 095) or bicycle (10 552) helmet. Jointly, the five risk factors accounted for 61% (528 105) of the deaths; this proportion is less than the sum of the five individual proportions, because many persons were at risk from more than one factor (further details are in online




Figure 2 Trends in death rates. The top line in each panel shows the observed rates (square) and the fitted line for all crash deaths of people potentially exposed to each risk factor. Each panel shows the observed rates (×) and fitted line for deaths attributed to each risk factor and for other deaths not attributed to that factor (triangle). The y axis uses a log scale so that convergence or divergence of changes in rates can be easily judged. Pointwise 95% confidence intervals are shaded.

Table 3 Rates (per 100 000 person-years) and change in rates, for all deaths, deaths attributed to each risk factor, and other deaths not attributed to each risk factor.*

Risk factor	Rates				
	1982†	2001†	Difference‡	Change (%)§	95% CI
Alcohol					
All deaths	19.57	14.39	-5.18	-26	-30 to -23
Attributed deaths	10.27	4.88	-5.39	-53	-56 to -49
Other deaths	9.52	9.67	+0.16	+2	-6 to +9
Not wearing a seat belt					
All deaths	13.62	11.53	-2.10	-15	-20 to -11
Attributed deaths	7.05	3.57	-3.48	-49	-52 to -47
Other deaths	6.77	8.11	+1.34	+20	+9 to +32
No air bag					
All deaths	12.11	10.04	-2.07	-17	-22 to -12
Attributed deaths	0.67	0.56	-0.11	-17	-38 to +12
Other deaths	11.44	9.48	-1.95	-17	-21 to -13
No motorcycle helmet					
All deaths	1.99	0.74	-1.24	-63	-72 to -51
Attributed deaths	0.43	0.11	-0.32	-74	-81 to -63
Other deaths	1.55	0.63	-0.93	-60	-69 to -47
No bicycle helmet					
All deaths	0.40	0.26	-0.14	-35	-40 to -30
Attributed deaths	0.26	0.16	-0.10	-39	-45 to -33
Other deaths	0.14	0.10	-0.04	-28	-32 to -23
All five risk factors					
All deaths	19.57	14.39	-5.18	-26	-30 to -23
Attributed deaths	14.22	7.36	-6.85	-48	-51 to -45
Other deaths	5.64	7.26	+1.62	+29	+18 to +41

*Not all people were considered exposed to all risk factors. For example, only vehicle occupants (not motorcyclists) were considered to be at risk for not wearing a seat belt and only motorcycle riders were considered to be potential wearers of motorcycle helmets. Death rates attributed to each risk factor alone sum to more than rates attributed to all five risk factors, as some deaths can be attributed to more than one factor.

†Rates for 1982 and 2001 are from the fitted models.

‡Rate difference = 2001 rate minus 1982 rate.

§Rate change = (2001 rate minus 1982 rate)/(1982 rate).

Appendices 2, 3, and 4; see <http://www.injuryprevention.com/supplemental>).

Over the 20 years, 46% of the dead were exposed to at least some alcohol; either they were a driver, pedestrian, or bicyclist with some level of blood alcohol, or their driver or another driver in the crash had some alcohol in their blood. Among dead vehicle occupants 75% wore no seat belt and 88% of front seat occupants had no air bag. Half (51%) of dead motorcyclists and 98% of dead bicyclists were not wearing a helmet.

All risk factors became less common among the dead over time (table 2, fig 1), but this change in prevalence was greatest for lack of an air bag (48% decrease from 1992 to 2001), and least for not wearing a bicycle helmet (4% decrease during 1994 to 2001). Prevalence of exposure to any alcohol fell 20% and not wearing a seat belt declined 39%.

Changes in death rates over time

Death rates attributed to alcohol decreased 53%, while the rate of other deaths not attributed to alcohol increased 2%,




Figure 3 Trends in death rates for motorcycle riders and bicycle riders. The top line in each panel shows the observed rates (square) and the fitted line for all crash deaths of riders. Each panel shows the observed (x) and fitted lines for deaths attributed to each risk factor and for other deaths not attributed to that factor (triangle). The y axis uses a log scale so that convergence or divergence of changes in rates can be easily judged. Pointwise 95% confidence intervals are shaded.

and the rate of all deaths decreased 26% (table 3 and fig 2). Among vehicle occupants, death rates attributed to not wearing a seat belt decreased 49%, while the rate of other deaths increased 20%; in combination these divergent trends produced a 15% decrease in crash death rates for all vehicle occupants.

For vehicle front occupants, death rates attributed to no air bag fell 17%, rates unrelated to an air bag decreased by 17%, and all deaths declined 17% (table 3, fig 2). When we limited the change estimates for air bags to 1992 through 2001, the interval during which air bag prevalence changed substantially, deaths attributed to no air bag changed by -42% (95% CI -51 to -32), other deaths changed by -1% (95% CI -5 to +2), and all front occupant deaths changed by -4% (95% CI -8 to 0).

Rates of deaths attributed to not wearing a motorcycle helmet fell 74%, other motorcycle death rates fell 60%, and overall motorcycle death rates declined 63% (table 3, fig 3). Observed rates of motorcycle deaths increased after 1997.

Rates of deaths attributed to not wearing a bicycle helmet decreased 39%, rates of other bicycle deaths fell 28%, and all bicycle death rates decreased 35% (table 3, fig 3). During 1994–2001, the years in which bicycle helmet prevalence changed, the change in attributed deaths was -24% (95% CI -30 to -17), the change in other deaths was -14% (95% CI -22 to -5), and the change in all bicycle deaths was -20% (95% CI -26 to -14).

For all five risk factors combined the rate of attributed deaths fell 48%, the rate of other deaths increased 29%, and the rate of all deaths decreased 26% (table 3).

Prevented deaths

Because changes in attributed death rates were always more favorable than changes in other death rates, the changes in risk factor prevalence resulted in positive estimates of prevented deaths (lives saved): during 1982–2001, 153 168 lives were saved by decreased drinking and driving, 129 297 by increased seat belt use, 4305 by increased air bag prevalence, 6475 by increased use of motorcycle helmets, and 239 by increased use of bicycle helmets. Jointly the favorable trends in all five risk factors prevented 293 898 deaths.

DISCUSSION

Of 858 741 deaths during the 20 years, 43% were attributed solely to alcohol use, 30% to not wearing a seat belt only, 4% to lack of an air bag, and 2.6% to not wearing a motorcycle or bicycle helmet. Jointly the five risk factors accounted for 61% of the deaths. These are estimates of the proportion of deaths that could have been prevented if each of the five study risk factors had been removed independently or all were removed simultaneously.

The interpretation of changes in attributed death rates is more difficult, as changes can occur for reasons unrelated to the studied risk factors; for example, changes in miles driven per person will change attributed death rates even if risk factor prevalence does not change. Therefore changes in attributed rates can be better interpreted in conjunction with information about changes in prevalence and changes in the rate of deaths not attributed to each risk factor.

The prevalence of some exposure to anyone's use of alcohol fell 20% among all who died in crashes during 1982–2001. Several studies have reported that alcohol use is strongly associated with crash risk^{14 35–42} and alcohol use can affect the risk of a crash for everyone on public roads. As a consequence, decreased use of alcohol alone made a larger contribution to the decline in crash death rates than any of the other four risk factors that we studied; an estimated 153 168 lives were saved by reduced alcohol use.

Not wearing a seat belt was second in importance. The association of no seat belt with death is not as strong as for

most observed alcohol blood levels and not wearing seat belts only affects people in vehicles. Nevertheless, the prevalence of this risk factor fell 39%, consistent with observed seat belt use,⁹ and an estimated 129 297 lives were saved.

Lack of an air bag showed the greatest decrease in prevalence (48%) of the factors we studied. But the contribution of this change was relatively small (4305 lives saved), as lack of an air bag is only weakly associated with the risk of death^{10 43–46} and air bags only affect front seat occupants.

Failure to wear a motorcycle helmet decreased somewhat and deaths attributed to this risk factor declined. But motorcycle death rates declined during this interval in part because there were fewer motorcycles. Registered motorcycles fell from 5.7 million in 1982 to 3.8 million in 1997; after 1997 registrations increased, reaching 4.9 million in 2001.^{1 47} Changes in motorcycle helmet use saved approximately 6475 lives.

Bicycle helmet use was uncommon during 1982–2001 and only 239 deaths were prevented by changes in helmet use.

Limitations

Counts of traffic crash deaths in FARS and in national death certificate data agree closely.⁴⁸ However blood alcohol levels were missing for over half the drivers, pedestrians, and bicyclists in this study. Several studies of alcohol in crashes have relied on singly imputed alcohol values, which ignore uncertainty regarding the missing values.^{49–51} Restricting an analysis to records with complete data leads to valid estimates only if data are missing completely at random,³⁴ which is not the case in crash data.³² We used multiple imputation, which relies on the weaker assumption that data were missing at random, conditional on the values of variables used for imputation.^{30 34} Bias could still be present if the pattern of missing values were strongly related to unmeasured aspects of crashes or people.

Any study of attributed deaths relies on assumptions about what fraction of deaths should be attributed to a risk factor. We used attributable fraction methods based on adjusted risk ratios.^{53 54} Several studies provide evidence that the risk ratio for a serious crash increases steeply with blood alcohol level of a driver, pedestrian, or bicyclist.^{14 35–42} Our estimates are relatively insensitive to choice of alcohol risk ratio estimates, as once the risk ratio is 10 or greater, the attributable fraction becomes 0.9 or more; 81% of the alcohol attributable fractions in our study were 0.9 or greater. The National Highway Traffic Safety Administration estimates that air bags reduce the risk of driver death by about 11%,⁵⁵ which differs little from the estimates we used. The agency estimates that seat belts reduce the risk of front occupant death by 45% in cars and 60% in light trucks.⁵⁶ Had we used these estimates, the number of deaths attributed to not wearing a seat belt would have been somewhat reduced.

Past studies

Previous studies of alcohol and driving, including reports using multiple imputation, have reported declines in alcohol related crash deaths in the US.^{50 57} These studies defined a crash as alcohol related if a driver, pedestrian, or bicyclist either had evidence of any drinking or had a level of blood alcohol above a certain level, such as 0.08 g/dl. Using this method, these studies have also reported decreases in deaths related to alcohol use.

Trends in crash deaths due to factors not studied

While the rate of death attributed jointly to the five risk factors decreased by 48%, the rate of other deaths not attributed to these factors increased by 29%. This suggests that favorable changes in the prevalence of the risk factors we studied exerted a downward influence on crash mortality,

Key points

- In the US during 1982–2001, 858 741 people died in traffic crashes.
- These deaths were attributed to one of five risk factors:
 - Alcohol use while driving (43%)
 - Failure to wear a seat belt (30%)
 - Lack of an air bag (4%)
 - Failure to wear a motorcycle helmet (1%)
 - Failure to wear a bicycle helmet (1%).
- 61% of the deaths were attributed to the joint effects of the five risk factors.
- The five risk factors became less common over the 20 year period. This favorable change saved lives:
 - Reduced use of alcohol—153 168 lives saved
 - Increased seat belt use—129 297 lives saved
 - Increased air bag prevalence—4305 lives saved
 - Increased use of motorcycle helmets—6475 lives saved
 - Increased use of bicycle helmets—239 lives saved.

while other changes increased crash mortality. One factor that may have increased crash mortality is faster driving speed; in 1987 many states raised the speed limit on rural freeways from 55 to 65 miles per hour⁵⁸ and after 1995 speed limits on many freeways were increased further.⁶ Another factor is the increase in miles driven, which increased from 6885 per person-year in 1982 to 9803 in 2001.¹ A third factor may be the shift in the vehicle fleet from passenger cars to light pickups and sport utility vehicles.^{6, 59}

CONCLUSION

Decreased drinking of alcohol and increased use of seat belts were associated with substantial reductions in crash mortality from 1982 through 2001. Increased presence of air bags, motorcycle helmets, and bicycle helmets also prevented some deaths. Favorable changes in all five risk factors were estimated to have changed the trend in death rates from a projected 29% rise to an observed 26% decline.

Authors' affiliations

P Cummings, Harborview Injury Prevention and Research Center and the Department of Epidemiology, University of Washington, Seattle, WA, USA

F P Rivara, Harborview Injury Prevention and Research Center, the Department of Pediatrics, and the Department of Epidemiology, University of Washington, WA, USA

C M Olson, Harborview Injury Prevention and Research Center, the Department of Medicine, and the Department of Epidemiology, University of Washington, Seattle, WA, USA

K M Smith, Harborview Injury Prevention and Research Center and the Department of Pediatrics, University of Washington, Seattle, WA, USA

Funding/support: This work was supported by grants R49/CCR002570 and R49/CCR019477-01 from The Centers for Disease Control and Prevention, Atlanta, Ga and by the Crash Injury Research and Engineering Network of the National Highway Traffic Safety Administration. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of The Centers for Disease Control and Prevention or the National Highway Traffic Safety Administration.

Competing interests: none.

REFERENCES

- 1 **National Highway Traffic Safety Administration.** Traffic Safety Facts 2002: A Compilation of Motor Vehicle Crash Data from the Fatality Analysis Reporting System and the General Estimates System. DOT HS 809 620. Washington, DC: National Highway Traffic Safety Administration, 2004.
- 2 **National Highway Traffic Safety Administration.** Traffic Safety Facts 2003: A Compilation of Motor Vehicle Crash Data from the Fatality Analysis Reporting System and the General Estimates System. DOT HS 809 775. Washington, DC: National Highway Traffic Safety Administration, 2005.
- 3 **Waller PF.** Challenges in motor vehicles safety. *Annu Rev Pub Health* 2002;**23**:93–113.
- 4 **Shults RA, Elder RW, Sleet DA, et al.** Reviews of evidence regarding interventions to reduce alcohol-impaired driving. *Am J Prev Med* 2001;**21**:66–88.
- 5 **Villaveces A, Cummings P, Koepsell TD, et al.** The association of alcohol-related laws with deaths due to motor vehicle and motorcycle crashes in the United States, 1980–1997. *Am J Epidemiol* 2003;**157**:131–40.
- 6 **Evans L.** *Traffic safety.* Bloomfield Hills, MI: Science Serving Society, 2004, 91–94, 211–216, 252–264, 295–298, 404.
- 7 **Rivara FP, Thompson DC, Cummings P.** Effectiveness of primary and secondary enforced seat belt laws. *Am J Prev Med* 1999;**16**(Suppl 1):S30–S39.
- 8 **Dinh-Zarr TB, Sleet DA, Shults RA, et al.** Reviews of evidence regarding interventions to increase the use of safety belts. *Am J Prev Med* 2001;**21**:48–65.
- 9 **Glassbrenner D.** Safety belt use in 2004—demographic results. DOT HS 809 848. Washington, DC: National Highway Traffic Safety Administration, National Center for Statistics and Analysis, 2005.
- 10 **Kahane CJ.** Fatality reduction by air bags: analysis of accident data through early 1996. DOT HS 808 470. Washington, DC: National Highway Traffic Safety Administration, 1996.
- 11 **Glassbrenner D.** Motorcycle helmet use in 2004—overall results. DOT HS 809 867. Washington, DC: National Highway Traffic Safety Administration, 2005.
- 12 **Rivara FP, Thompson DC, Patterson MQ, et al.** Prevention of bicycle-related injuries: helmets, education, and legislation. *Ann Rev Public Health* 1998;**19**:293–318.
- 13 **MacMahon B, Trichopoulos D.** *Epidemiology: principles and methods*, 2nd ed. Boston: Little, Brown, 1996:291–9.
- 14 **Zador PL, Krawchuk SA, Voas RB.** Alcohol-related relative risk of driver fatalities and driver involvement in fatal crashes in relation to driver age and sex: an update using 1996 data. Washington, DC: National Highway Traffic Safety Administration, 1999.
- 15 **Olson CM, Cummings P, Rivara FP.** The association of first and second generation air bags with front occupant death in car crashes: a matched cohort study. *Am J Epidemiol* (in press).
- 16 **Norvell DC, Cummings P.** Association of helmet use with death in motorcycle crashes: a matched-pair cohort study. *Am J Epidemiol* 2002;**156**:483–7.
- 17 **Thompson DC, Rivara FP, Thompson RS.** Effectiveness of bicycle helmets in preventing head injuries: a case-control study. *JAMA* 1996;**276**:1968–73.
- 18 **Lumley T, Diehr P, Emerson S, et al.** The importance of the normality assumption in large public health data sets. *Annu Rev Public Health* 2002;**23**:151–69.
- 19 **US Census Bureau.** Quarterly population estimates. Available at http://www.census.gov/popest/archives/1980s/80s_nat_detail.html (accessed January 2005).
- 20 **Breslow NE, Day NE.** *Statistical methods in cancer research. Volume II - the design and analysis of cohort studies.* Lyon, France: International Agency for Research on Cancer, 1987:120–76.
- 21 **Cummings P, Norton R, Koepsell TD.** Rates, rate denominators, and rate comparisons. In: Rivara FP, Cummings P, Koepsell TD, Grossman DC, Maier RV (eds). *Injury control: a guide to research and program evaluation.* New York, NY: Cambridge University Press, 2001:64–74.
- 22 **Vindicator 2001 - Release No 1.** Arlington, VA: Highway Loss Data Institute, 2001.
- 23 **Raghuathan TE.** What do we do with missing data? Some options for analysis of incomplete data. *Annu Rev Public Health* 2004;**25**:99–117.
- 24 **Schimert J, Schafer JL, Hesterberg T, et al.** *Analyzing data with missing values in S-Plus.* Seattle, WA: Insightful Corporation, 2001.
- 25 **S-PLUS 6.2 for Windows, Professional Edition.** Seattle, WA: Insightful Corporation, 2003.
- 26 **van Buuren S, Boshuizen HC, Knook DL.** Multiple imputation of missing blood pressure covariates in survival analysis. *Stat Med* 1999;**18**:681–94.
- 27 **Stata Statistical Software: Release 8.2.** College Station, TX: Stata Corporation, 2004.
- 28 **Royston P.** Multiple imputation of missing values. *Stata J* 2004;**4**:227–41.
- 29 **Rubin DB, Schafer JL, Subramanian R.** Multiple imputation of missing blood alcohol concentration (BAC) values in FARS. DOT HS 808 816. Washington DC: National Highway Traffic Safety Administration, 1998.
- 30 **Subramanian R.** Transitioning to multiple imputation—a new method to estimate missing blood alcohol concentration (BAC) values in FARS. DOT HS 809 403. Washington, DC: National Highway Traffic Safety Administration, 2002.
- 31 **Harborview Injury Prevention and Research Center.** Data on alcohol, seatbelts and air bags added to FARS. Available at <http://depts.washington.edu/hiprc/FARS/FARSdefine.html> (accessed March 2005).
- 32 **Wooldridge JM.** *Econometric analysis of cross section and panel data.* Cambridge, MA: The MIT Press, 2002:649–50.

- 33 **Greenland S.** Model-based estimation of relative risks and other epidemiologic measures in studies of common outcomes and in case-control studies. *Am J Epidemiol* 2004;**160**:301–5.
- 34 **Schafer JL.** *Analysis of incomplete multivariate data.* New York: Chapman & Hall, 1997, 1–2, 108–10, 257–75.
- 35 **Holcomb RL.** Alcohol in relation to traffic accidents. *J Am Med Assoc* 1938;**111**:1076–85.
- 36 **Haddon W Jr,** Valien P, McCarrroll JR, et al. A controlled investigation of the characteristics of adult pedestrians fatally injured by motor vehicles in Manhattan. *J Chron Dis* 1961;**14**:655–78.
- 37 **McCarrroll JR,** Haddon W Jr. A controlled study of fatal automobile accidents in New York City. *J Chron Dis* 1962;**15**:811–26.
- 38 **Borkenstein RF,** Crowther RF, Shumate RP, et al. *The role of the drinking driver in traffic accidents.* Bloomington, Indiana: Indiana University, Department of Police Administration, 1964:1–245.
- 39 **Borkenstein RF,** Crowther RF, Shumate RP, et al. The role of the drinking driver in traffic accidents (The Grand Rapids Study). *Blutalkohol* 1974;**11**(Suppl 1):1–132.
- 40 **Zador PL.** Alcohol-related relative risk of fatal driver injuries in relation to driver age and sex. *J Stud Alcohol* 1991;**52**:302–10.
- 41 **Zador PL,** Krawchuk SA, Voas RB. Alcohol-related relative risk of driver fatalities and driver involvement in fatal crashes in relation to driver age and gender: an update using 1996 data. *J Stud Alcohol* 2000;**61**:387–95.
- 42 **Li G,** Baker SP, Smialek JE, et al. Use of alcohol as a risk factor for bicycling injury. *JAMA* 2001;**285**:893–6.
- 43 **Ferguson SA,** Lund AK, Greene MA. *Driver fatalities in 1985–1994 air bag cars.* Arlington, VA: Insurance Institute for Highway Safety, 1995.
- 44 **Braver ER,** Ferguson SA, Greene MA, et al. Reductions in deaths in frontal crashes among right front passengers in vehicles equipped with passenger air bags. *JAMA* 1997;**278**:1437–9.
- 45 **Cummings P,** Koepsell TD, Rivara FP, et al. Air bags and passenger fatality according to passenger age and restraint use. *Epidemiology* 2002;**13**:525–32 [erratum 749].
- 46 **Cummings P,** McKnight B, Rivara FP, et al. Association of driver air bags with driver fatality: a matched cohort study. *BMJ* 2002;**324**:1119–22.
- 47 **Shankar U.** Recent trends in motorcycle crashes. DOT HS 809 271. Washington, DC: National Highway Traffic Safety Administration, 2001.
- 48 **Vyrostek SB,** Annett JL, Ryan GW. Surveillance for fatal and nonfatal injuries - United States, 2001. Surveillance Summaries. *MMWR Morb Mortal Wkly Rep* 2004;**53**:SS 1–57.
- 49 **Lund AK,** Wolfe AC. Changes in the incidence of alcohol-impaired driving in the United States, 1973–1986. *J Stud Alcohol* 1991;**52**:293–301.
- 50 **National Highway Traffic Safety Administration.** Alcohol involvement in fatal crashes 2000. DOT HS 809 419. Washington, DC: National Highway Traffic Safety Administration, 2002.
- 51 **Quinlan KP,** Brewer RD, Sleet DA, et al. Characteristics of child passenger deaths and injuries involving drinking drivers. *JAMA* 2000;**283**:2249–52.
- 52 **Klein TM.** A method for estimating posterior BAC distributions for persons involved in fatal traffic accidents. DOT HS 807 094. Washington, DC: National Highway Traffic Safety Administration, 1986.
- 53 **Benichou J.** A review of adjusted estimators of attributable risk. *Stat Methods Med Res* 2001;**10**:195–216.
- 54 **Gefeller O,** Land M, Eide GE. Averaging attributable fractions in the multifactorial situation: assumptions and interpretation. *J Clin Epidemiol* 1998;**51**:437–41.
- 55 **National Highway Traffic Safety Administration.** Fifth/Sixth Report to Congress: Effectiveness of occupant protection systems and their use. DOT HS 809 442. Washington, DC: National Highway Traffic Safety Administration, 2001.
- 56 **Kahane CJ.** Fatality reduction by safety belts for front-seat occupants of cars and light trucks: updated and expanded estimates based on 1986–99 FARS data. DOT HS 809 199. Washington, DC: National Highway Traffic Safety Administration, 2000.
- 57 **Subramanian R.** Alcohol involvement in fatal crashes 2001. DOT HS 809 579. Washington, DC: National Highway Traffic Safety Administration, National Center for Statistics and Analysis, 2003.
- 58 **Ossiander E,** Cummings P. Freeway speed limits and traffic fatalities in Washington State. *Accid Anal Prev* 2002;**34**:13–18.
- 59 **Rivara FP,** Cummings P, Mock C. Injuries and death of children in rollover motor vehicle crashes in the United States. *Inj Prev* 2003;**9**:76–80.

LACUNAE

What should drivers do with their whips?

The Highway Code sells one million copies a year in the UK, and everyone who has taken their driving test has read it. Trouble is, we tend to forget the best practices. This year, the Highway Code, first published in 1931, celebrates its 75th anniversary. This is what the Highway Code said...

1931 edition: What should drivers do with their whips? Drivers should "rotate the whip above the head, then incline the whip to the right or left to show the direction in which the turn is to be made." This, of course, applies to horse-drawn vehicles only.

If a driver flapped his wrist out of the window, what did this indicate? In those days, it was the hand signal for stopping: "Extend the right arm with the palm of the hand turned downwards and move the arm slowly up and down, keeping the wrist loose."

How did motorists indicate an intention to overtake? By beeping their horn.

1954 edition: What was indicated by the road sign that shows a flaming torch? Not an Olympic stadium or "arsonists at work". For many years, this was the sign indicating that a school was nearby.

What was the "wrong" side of the road? According to the 1954 Code, motorists were not allowed to "park at night on the 'wrong' side of the road". While the Code offers no guidance as to what the "wrong" side is, parking facing against the traffic direction was once discouraged.

What is the difference between the stopping times for 50mph in the 1954 edition and the present day? Actually, there isn't one. Despite all the changes in motoring technology during the previous half century, the answer then —175 feet—is exactly the same as now, except that now it is given in metric—53m—first.

2006 edition: If you are a walking along a pavement or footpath, what should you take care to do? The Code says that, where possible, avoid walking next to the kerb with your back to the traffic.

Can you take a mobile phone call while driving at 30mph in a built-up area? The Code says: "You must not use a hand-held mobile phone, or similar device, when driving or when supervising a learner driver."

Contributed by Ian Scott

Appendix 1

Risk ratios for death and attributable fractions associated with each study risk factor, compared with not having that risk factor

Driver, pedestrian, or bicyclist blood alcohol level				
Blood alcohol level (g/dl)				Attributable fraction
Blood alcohol level (g/dl)	Age (y)	Sex	Risk ratio	Attributable fraction
0	All	Both	1.00 (Reference)	0
0.01	0–20	Female	1.22	0.18
0.01	0–20	Male	1.42	0.30
0.01	≥21	Both	1.00	0
0.02–0.04	0–20	Female	1.98	0.49
0.02–0.04	0–20	Male	3.44	0.71
0.02–0.04	21–34	Both	2.04	0.51
0.02–0.04	≥35	Both	2.02	0.50
0.05–0.07	0–20	Female	3.56	0.72
0.05–0.07	0–20	Male	9.94	0.90
0.05–0.07	21–34	Both	3.76	0.73
0.05–0.07	≥35	Both	3.70	0.73
0.08–0.09	0–20	Female	5.80	0.83
0.08–0.09	0–20	Male	24.03	0.96
0.08–0.09	21–34	Both	6.25	0.84
0.08–0.09	≥35	Both	6.13	0.84
0.10–0.14	0–20	Female	11.50	0.91
0.10–0.14	0–20	Male	82.73	0.99

Driver, pedestrian, or bicyclist blood alcohol level

Blood alcohol				Attributable
level (g/dl)	Age (y)	Sex	Risk ratio	fraction
0.10–0.14	21–34	Both	12.74	0.92
0.10–0.14	≥35	Both	12.41	0.92
≥0.15	0–20	Female	73.62	0.99
≥0.15	0–20	Male	2371.74	1.00
≥0.15	21–34	Both	88.13	0.99
≥0.15	≥35	Both	84.13	0.99

Not wearing a seat belt

Seat belt	Seat position	Age (years)	Air bag present	Risk ratio	Attributable fraction
Yes	All	All	Yes or no	1.00 (Reference)	0
No	Front	0–49	No	2.33	0.57
No	Front	0–49	Yes	2.55	0.61
No	Front	≥50	No	1.90	0.47
No	Front	≥50	Yes	2.08	0.52
No	Rear	0–49	No	1.95	0.49
No	Rear	≥50	No	1.59	0.37

Not having a front seat airbag

Air bag	Seat position	Age (years)	Seat belt	Risk ratio	Attributable fraction
Yes	All front	All	Yes or	1.00 (Reference)	0

Driver, pedestrian, or bicyclist blood alcohol level

Blood alcohol					Attributable
level (g/dl)	Age (y)	Sex		Risk ratio	fraction
			No		
No	Driver	5–12	No	0.96	–0.04
No	Driver	5–12	Yes	1.05	0.05
No	Driver	13–49	No	1.03	0.03
No	Driver	13–49	Yes	1.12	0.11
No	Driver	≥50	No	1.06	0.06
No	Driver	≥50	Yes	1.16	0.14
No	Front	0–4	No	0.71	–0.41
	passenger				
No	Front	0–4	Yes	0.78	–0.28
	passenger				
No	Front	5–12	No	0.96	–0.04
	passenger				
No	Front	5–12	Yes	1.05	0.05
	passenger				
No	Front	13–49	No	1.06	0.06
	passenger				
No	Front	13–49	Yes	1.16	0.14
	passenger				
No	Front	≥50	No	1.20	0.17
	passenger				

Driver, pedestrian, or bicyclist blood alcohol level

Blood alcohol level (g/dl)				Risk ratio	Attributable fraction
No	Front passenger	≥50	Yes	1.31	0.24

No seat belt, or no air bag, or no seat belt and no airbag

Seat belt	Air bag	Seat position	Age (years)	Risk ratio	Attributable fraction
Yes	Yes	All	All	1.00 (Reference)	0
Yes	No	Driver	5–12	0.96	–0.04
Yes	No	Driver	13–49	1.03	0.03
Yes	No	Driver	≥50	1.06	0.06
Yes	No	Front passenger	0–4	0.71	–0.41
Yes	No	Front passenger	5–12	0.96	–0.04
Yes	No	Front passenger	13–49	1.06	0.06
Yes	No	Front passenger	≥50	1.20	0.17
No	Yes	Driver	0–49	2.33	0.57
No	Yes	Driver	≥50	1.90	0.47
No	Yes	Front passenger	0–49	2.33	0.57

Driver, pedestrian, or bicyclist blood alcohol level					
Blood alcohol level (g/dl)					Attributable fraction
Age (y)	Sex			Risk ratio	
No	Yes	Front passenger	≥50	1.90	0.47
No	No	Driver	5–12	2.45	0.59
No	No	Driver	13–49	2.62	0.62
No	No	Driver	≥50	2.21	0.55
No	No	Front passenger	0–4	1.82	0.45
No	No	Front passenger	5–12	2.45	0.59
No	No	Front passenger	13–49	2.69	0.63
No	No	Front passenger	≥50	2.49	0.60
No	No	Rear seat	0–49	1.95	0.49
No	No	Rear seat	≥50	1.59	0.37
No motorcycle helmet					
Motorcycle helmet	Seat position				
Yes	All			1.00 (Reference)	0
No	Driver			1.54	0.35
No	Rear			1.72	0.42

Driver, pedestrian, or bicyclist blood alcohol level

Blood alcohol Attributable

level (g/dl) Age (y) Sex Risk ratio fraction

No bicycle helmet

Bicycle helmet

Yes 1.00 (Reference) 0

No 2.86 0.65

Appendix 2

Traffic crash deaths attributed to alcohol by age and subgroup, US, 1982–2001

		All deaths			Deaths attributed to anyone's drinking alcohol			Deaths attributed to drinking alcohol by the dead person			Deaths attributed to drinking by someone other than the dead person		
Subgroup		Count	Rate	Count	Rate	Alcohol (%)†	Count	Rate	Alcohol (%)†	Count	Rate	Alcohol (%)†	
All		858,741	16.75	366,606	7.15	43	245,830	4.80	29	144,359	2.82	17	
Age group (years)	0–15	67,867	5.73	14,331	1.21	21	1482	0.13	2	13,066	1.10	19	
	16–20	130,933	34.39	57,446	15.09	44	32,019	8.41	24	28,210	7.41	22	
	21–30	210,018	25.92	122,395	15.11	58	87,720	10.83	42	42,943	5.30	20	
	31–64	321,283	15.18	151,541	7.16	47	113,482	5.36	35	49,303	2.33	15	
	65+	128,640	20.30	20,893	3.30	16	11,127	1.76	9	10,837	1.71	8	

	Subgroup	All deaths		Deaths attributed to anyone's drinking alcohol			Deaths attributed to drinking alcohol by the dead person			Deaths attributed to drinking by someone other than the dead person		
		Count	Rate	Count	Rate	Alcohol	Count	Rate	Alcohol	Count	Rate	Alcohol
						(%)†			(%)†			(%)†
Persons	Drivers	443,655	8.66	197,642	3.86	45	175,204	3.42	39	33,350	0.65	8
	Passengers	213,667	4.17	79,116	1.54	37	0	0	0	79,116	1.54	37
	Pedestrians	121,135	2.36	53,467	1.04	44	42,354	0.83	35	20,893	0.41	17
	Motorcyclists	63,811	1.24	31,514	0.61	49	25,297	0.49	40	8,393	0.16	13
	Bicyclists	16,473	0.32	4866	0.09	30	2975	0.06	18	2,606	0.05	16

*Driver and passenger categories exclude motorcyclists. Deaths, rates, and proportions attributed to drinking by each person and to drinking by another person sum to more than all alcohol deaths, rates, and proportions, because some deaths can be attributed to drinking by more than one person.

†Alcohol (%) = $100 \times \text{alcohol deaths/all deaths}$.

Rates per 100,000 person-years.*

Appendix 3

Traffic crash deaths attributed to not wearing a seat belt and not having an airbag: US data for vehicle occupants, 1982–2001

Age (years)	Deaths of vehicle occupants	Deaths attributed to no seat belt			Deaths of vehicle front occupants	Deaths attributed to no air bag*		
		Deaths				Deaths		
		Deaths	Rate	(%)†		Deaths	Rate	(%)†
All	641,114	259,239	5.06	40	564,124	31,377	0.61	6
0–15	37,530	14,443	1.22	38	18,455	–1337‡	–0.11‡	–7‡
16–20	106,395	48,061	12.62	45	92,970	4386	1.15	5
21–30	158,393	73,574	9.08	46	144,439	6325	0.78	4
31–64	239,574	96,330	4.55	40	216,542	12,617	0.60	6
65+	99,222	26,831	4.23	27	91,718	9387	1.48	10

*Airbag analysis limited to front occupants in vehicles of the type that customarily have air bags.

†Deaths (%) = attributed deaths/all deaths.

‡Negative attributed deaths are lives saved among the young by not being exposed to air bags, as air bags were a net hazard to young children. Rates per 100,000 person-years.

Appendix 4

Traffic crash deaths attributed to not wearing a helmet among motorcycle riders and bicyclists by age: US data, 1982–2001

Age (years)	Deaths of motorcycle riders	Deaths attributed to no motorcycle helmet			Deaths of bicyclists	Deaths attributed to no bicycle helmet		
		Deaths		Deaths (%)*		Deaths		Deaths (%)*
		Deaths	Rate			Deaths	Rate	
All	63,811	12,095	0.24	19	16,473	10,552	0.21	64
0–15	1,896	434	0.04	23	6523	4210	0.36	65
16–20	11,016	2225	0.58	20	1744	1125	0.30	65
21–30	26,356	5040	0.62	19	2264	1455	0.18	64
31–64	23,608	4258	0.20	18	4817	3045	0.14	63
65+	935	138	0.02	15	1125	717	0.11	64

*Deaths (%) = Attributed deaths/all deaths.
Rates per 100,000 person-years.