



Triangulating case-finding tools for patient safety surveillance: a cross-sectional case study of puncture/laceration

Jennifer A Taylor,¹ Daniel Gerwin,¹ Laura Morlock,² Marlene R Miller³

► Additional appendices are published online only. To view these files please visit the journal online (<http://injuryprevention.bmj.com>).

¹Department of Environmental and Occupational Health, Drexel University School of Public Health, Philadelphia, Pennsylvania, USA

²Department of Health Policy and Management, The Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

³Department of Pediatrics, Johns Hopkins Children's Center and Department of Health Policy and Management, The Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

Correspondence to

Dr Jennifer Taylor, Department of Environmental and Occupational Health, Drexel University School of Public Health, 1505 Race Street, MS 1034, Philadelphia, PA 19102, USA; Jat65@drexel.edu

Accepted 16 March 2011
Published Online First
5 May 2011

ABSTRACT

Objective To evaluate the need for triangulating case-finding tools in patient safety surveillance. This study applied four case-finding tools to error-associated patient safety events to identify and characterise the spectrum of events captured by these tools, using *puncture* or *laceration* as an example for in-depth analysis.

Data sources/study setting Retrospective hospital discharge data were collected for calendar year 2005 (n=48 418) from a large, urban medical centre in the USA.

Study design The study design was cross-sectional and used data linkage to identify the cases captured by each of four case-finding tools.

Data collection/extraction methods Three case-finding tools (International Classification of Diseases external (E) and nature (N) of injury codes, Patient Safety Indicators (PSI)) were applied to the administrative discharge data to identify potential patient safety events. The fourth tool was Patient Safety Net, a web-based voluntary patient safety event reporting system.

Results The degree of mutual exclusion among detection methods was substantial. For example, when linking *puncture* or *laceration* on unique identifiers, out of 447 potential events, 118 were identical between PSI and E-codes, 152 were identical between N-codes and E-codes and 188 were identical between PSI and N-codes. Only 100 events that were identified by PSI, E-codes and N-codes were identical. Triangulation of multiple tools through data linkage captures potential patient safety events most comprehensively.

Conclusions Existing detection tools target patient safety domains differently, and consequently capture different occurrences, necessitating the integration of data from a combination of tools to fully estimate the total burden.

BACKGROUND

In March 2000, the Institute of Medicine released its seminal public health report, 'To Err is Human: Building a Safer Health Care System', citing an annual estimate of 44 000–98 000 deaths in the USA due to medical errors.¹ Debate ensued about the estimates, and the argument as to how many events occur annually has yet to be resolved.^{2–3} The Institute of Medicine report provided the first US medical error mortality estimates, but a surveillance system to consistently capture the full spectrum of patient safety events has not been developed.

A surveillance approach to patient safety is visualised (figure 1). Medical injuries are united with medical errors under the umbrella of surveillance,

allowing a more thorough description of the burden of patient safety events. Different case-finding tools have evolved to identify and measure patient safety events, and the role of each can be seen (table 1).

Medical record coding

Administrative data from inpatient hospital discharges capture *injuries not associated with error* and *error-associated injuries*. Within hospitals, nosologists review patient charts and assign codes for reimbursement purposes. These codes emanate from the International Classification of Diseases (ICD-9-CM), and relevant codes include 'nature of injury' (N-code) and 'external cause of injury' (E-code). Researchers have analysed administrative data coding to identify potential patient safety events in Victoria, Australia, and in American state-wide studies in Utah, Missouri and Wisconsin.^{4–7}


Hospital discharge data before 2008 are vulnerable to false positives, resulting from a failure to distinguish conditions present on admission (POA) from those acquired during hospitalisation. In 2008, the Centers for Medicare and Medicaid Services required the assignment of a POA code to every diagnosis, identifying whether or not the condition was acquired in hospital. Houchens *et al* report that hospitals have been found to over-report POA, probably because this practice reduces the apparent rate of injuries associated with error.⁸ POA coding has yet to be fully validated.

Administrative data may have limited usefulness owing to a lack of clinical detail or contextual information, variability in coding across providers, incentives for 'upcoding' or 'downcoding' and inaccuracy. Iezzoni states that 'reimbursement and reporting incentives can significantly shift coding practices'.⁹ Administrative data are nonetheless critical, because they are routinely collected, easily acquired and standardised for international comparison.

The patient safety indicators

The Patient Safety Indicators (PSI) are an administrative data algorithm, developed by the Agency for Healthcare Research and Quality (AHRQ) as screening indicators for patient safety case-finding and performance improvement at the provider and system level in hospitals.¹⁰ The PSI algorithm uses variables such as ICD-9-CM coded diagnoses and procedures, age, gender, admission source and discharge status to reflect preventable injury exposure more precisely. The principal difference between PSI and E- and N-codes is that PSI target only 20 in-hospital *error-associated injuries*,

Figure 1 Conceptual diagram of patient safety event classifications with relevant data sources and case-finding tools.



restricting numerators and denominators solely to events deemed preventable, by excluding patients whose pre-existing conditions would probably lead to PSI events regardless of error. There is a small amount of overlap between the ICD-9-CM codes used in the PSI categories and the adverse event range of N- and E-codes used in this study: eight use a few N-codes for adverse events (PSI 4, 7, 9, 10, 13, 15, 16) and five use a limited number of E-codes (postoperative hip fracture, birth trauma, foreign body, transfusion reactions and puncture/laceration).¹¹ A number of PSI have been found to incorporate false positives contained in the administrative data owing to the inclusion of pre-existing conditions.^{12 13} With respect to their narrowly defined outcomes, PSI have been found to have high specificity, moderate sensitivity and moderate predictive positive value, though these metrics vary considerably among the different PSI.^{11 14}

Voluntary error reporting

Voluntary error reporting captures a vast spectrum of patient safety events.¹⁵ Low reporting rates are the norm, and comparisons

with chart review have found that voluntary reports identified only 1.5% of adverse events and 6% of adverse drug events. The small, non-random fraction of incidents that is reported cannot be used to establish a numerator when calculating error rates or broad estimates.¹⁶

STUDY GOAL

Our objective was to examine the extent to which a fuller spectrum of *error-associated injuries* might be identified using four case-finding tools (E-codes, N-codes, PSI and Patient Safety Net (PSN)). We hypothesised that because each case-finding tool is differently designed and targeted, there would be minimal overlap in events identified by each tool. In so far as case-finding tools did not identify the same events, triangulation would be needed to achieve more comprehensive surveillance. We define triangulation as the use of data linkage to integrate the different results produced by applying multiple case-finding tools to the same patient cohort, using unique identifiers to remove

Table 1 Strengths and weaknesses of case-finding tools

Tool	Purpose	Key strengths	Key weaknesses
ICD-9-CM External cause of injury (E-codes)	Injury Surveillance and prevention	Standardised for international comparison; captures adverse drug reactions	Inconsistently used among hospitals; lack of clinical detail/context; variability in coding; inaccuracy
ICD-9-CM Nature of injury (N-codes)	Reimbursement	Routinely collected by all hospitals; easily acquired; standardised for international comparison	Lack of clinical detail/context; variability in coding; financial pressures for 'upcoding' or 'downcoding'; inaccuracy
Patient Safety Indicators (PSI)	Screening hospital discharge data for patient safety events to improve provider performance	More precisely reflects exposure by restricting the denominator to exclude patients whose pre-existing conditions would probably lead to PSI events regardless of error; international utility	Includes only 20 narrowly defined patient safety outcomes; does not capture falls or adverse drug reactions
Voluntary reporting (PSN)	Identifying medical errors	Encompasses all possible errors and near misses; captures both falls and medication errors	Low reporting rate; non-representative sample

PSN, Patient Safety Net.

duplicate events. We conducted a more in-depth investigation of *puncture/laceration* as a case study.

STUDY DESIGN

The study design was cross-sectional and used retrospective data. The setting was a large, urban medical centre. In 2005, the hospital had approximately 1900 total beds and 48 000 discharges. This study used two data sources: the hospital's inpatient discharge data for 2005 and the voluntary error-reporting system for the 2005 calendar year, the first whole year during which the voluntary error-reporting system was operational at the medical centre. All study activities were approved by the institutional review board at the investigators' institution and HIPAA (Health Information Portability and Accountability Act) approval was given by the participating hospital.

Case finding tools

External cause and nature of injury codes

The hospital's discharge data contain 15 diagnosis fields. The N-code and E-code criteria were chosen from the ICD-9-CM categories for adverse events (online supplementary appendix). This study grouped potential adverse events according to the following E-code categories: 'Misadventures to patient during surgical and medical care' (E870–E876), 'Surgical and medical procedures as the cause of abnormal reaction of patients or later complication, without mention of misadventure at the time of procedure' (E878–E879) and 'Drugs, medicinal and biological substances causing adverse effects in therapeutic use' (E930–E949). We included E930–E949 because there is evidence to suggest that a significant percentage of adverse drug events are due to medical errors which can be prevented through improvements in hospital systems.¹⁷ We used E850–E858 'Accidental poisoning by drugs, medical substances and biological substances' because this range included further opportunities to capture errors in drug administration. Finally, we included N-codes in the range of 960–979 'Poisonings by drugs, medicinal and biological substances', 995 'Certain adverse effects not elsewhere classified' and 996–999 'Complications of surgical and medical care not elsewhere classified'. We used these codes exactly as they appeared in the data because we were examining them as they would be used for surveillance purposes, rather than conducting a validation study.

Patient Safety Indicators

AHRQ's publicly available Windows QI software (version 3.0) was applied to the hospital's administrative discharge data.¹⁰ The algorithm assigned one of the 20 PSI categories to each discharge that met the criteria for a potential patient safety event.

Patient Safety Net

Patient Safety Net version 3.0 (PSN) is a voluntary, electronic error-reporting tool developed by the University HealthSystems Consortium (<https://www.uhc.edu/11851.htm>). Providers enter events into a web-based computer system through which each error is classified by event type and scored to indicate the extent of harm to the patient (A–I: A=unsafe conditions, I=death). Cases were selected that had a harm score of 'C' or higher, indicating that an event had reached the patient.

METHODS

Given the variable sensitivity and specificity of our case-finding tools, and because our study design did not include chart review, we do not refer to patient safety events in our study, but rather

to *potential patient safety events*. In order to generate patient-level summary statistics, a dichotomous variable was created for each of the four case identification criteria within a patient encounter: any discharge identified as having a potential patient safety event was assigned a numerical value of '1'. Non-events were given a value of '0'. These tabulations were not based on the first detection of an event by only one case-finding tool, but rather included any discharge as having an event if it was detected by PSN, PSI, N-code or E-code. Data were managed in Microsoft Access. The study employed the STATA software package (Stata 8, StataCorp LP) to perform all analyses.

Differences between potential events and non-events were summarised by age, length of stay, charges, discharge disposition, gender and complexity. Complexity was assessed through the APRDRG Patient Complexity Level (3M APRDRG grouping software, Ver 12). (<http://www.hcup-us.ahrq.gov>) Each patient was assigned an APRDRG value: 1 (low) to 4 (high). For the analysis, APRDRG complexity was dichotomised into high (3 or 4) and low (1 or 2) categories.

To account for similarities among categories in the different case identification tools, a crosswalk was created showing agreement of categories. For example, E-code 870, '*accidental cut, puncture, perforation, or haemorrhage during medical care*' (hereafter referred to as '*puncture/laceration*'), was determined to identify the same injury mechanism as PSI 15, '*accidental puncture/laceration*'. Of the 20 PSI categories, only three types of patient safety events had classifications in each of the other three case-finding tools: *foreign body*, *transfusion reactions* and *puncture/laceration*. As there were only three outcomes that could potentially be captured by all four case-finding tools, we examined additional patient safety outcomes from among PSN's event-type categories with classifications in two or more case-finding tools (table 2). We chose PSN as the template for comparison because PSN's categories are the most comprehensive, and include patient safety events of high interest, such as falls and medication errors.

In order to ascertain the number of identical events captured by multiple case-finding tools (overlap), data linkage was conducted. Potential events were linked on their encounter number for the hospital admission. Medical record number, name and demographic data were used to ensure valid data linkage. Total event counts from each case-finding tool were tabulated, and the overlap was identified.

For our case study of *puncture/laceration*, we took an additional step developed by Gallagher *et al*¹¹ to eliminate those cases in which *puncture/laceration* was POA. To achieve this, we removed events in which the patient had *puncture/laceration* as both the primary and secondary diagnosis. Since chart review was outside the scope of our study, this additional filtering process was an important validation method, an approach also used in a comparable study by Naessens *et al* in 2009.¹⁸

RESULTS

During the 2005 calendar year, 48 418 discharges were recorded. Consistent with previous findings, inpatient encounters flagged by at least one case-finding tool were more likely to have higher average total charges and higher average lengths of stay.^{19–21} Patients who had potential patient safety events were older and had higher levels of complexity. Some of the case-finding tools detected the same event, therefore an overestimate of patient safety events is presented (40%, table 3). For example, 78% of events detected by N-codes and E-codes are for the same event. De-duplication was performed only on the subset of the events in the crosswalk (table 2).

Table 2 Crosswalk of all case-finding criteria showing overlapping cases

PSN event type	Case-finding method				Overlapping events (N)
	PSN N	PSI N	N-code N	E-code N	
A. Medication error (ICD-9-CM: E873, E850-858)	1035	—	—	29	3
B. Adverse drug reaction (not a medication error) (ICD-9-CM: 960-979, E930-E949)	13	—	768	2501	0
C. Equipment/supplies (ICD-9-CM: 996, E874, E976.2)	98	—	5021	2	0
D. Fall (PSI 8)	373	3	—	—	2
E. Error related to procedure/treatment/test	1234	—	—	—	—
<i>Unintended laceration or puncture (PSI 15; ICD-9-CM: 998.2, E870)</i>	2	328	321	205	100
<i>Foreign body in patient (PSI 5; ICD-9-CM: 998.4, E871)</i>	0	3	2	8	0
F. Complication of procedure/treatment/test	220	—	—	—	—
<i>Wound dehiscence (PSI 14)</i>	1	1	—	—	0
<i>Pulmonary embolism (PSI 12)</i>	1	298	—	—	0
<i>Anaesthesia events (PSI 1)</i>	2	16	—	—	0
<i>Birth trauma (PSI 17)</i>	1	18	—	—	0
G. Transfusion	32	—	—	—	—
Transfusion reaction (PSI 16; ICD-9-CM: 999, E876.0)	3	4	8	0	0
H. Behavioural	40	—	—	—	—
I. Skin integrity	129	—	—	—	—
<i>Pressure ulcer (PSI 3)</i>	7	165	—	—	0
J. Care coordination/records	87	—	—	—	—
K. Other/miscellaneous	335	—	—	—	—
Total	3613	836	6120	2745	105

Italics indicate subcategories of PSN in common with one or more case-finding methods.

Parentheses indicate common categories among case-finding tools.

E-code, ICD-9-CM external cause of injury; N-code, ICD-9-CM nature of injury; PSI, Patient Safety Indicators; PSN, Patient Safety Net.

— indicates no category.

See appendices for complete results from each case-finding tool.

A summary of case-finding results is provided (table 4). The hospital's overall E-coding rate was 84%, indicating the percentage of discharges in which an E-code was assigned given that an N-code (ICD-9-CM 800-999) also appeared in the diagnostic string. Detailed tabulations of all events detected by each case-finding tool are available (online appendix).

The number of potential patient safety events flagged by each of the case-finding tools indicated that minimal overlap was

found after data linkage (table 2). *Puncture/laceration* provides an example. Applying the procedure of Gallagher *et al* eliminated 13 out of 205 events identified by E-codes, and 16 out of 321 events that were identified by N-codes (10 of these were the same events, identified by both E and N-codes). The total number of *puncture/lacerations* was thus reduced from 476 to 447, a 6% reduction, which is comparable to the 5% reduction found by Gallagher *et al*.¹¹

Table 3 Summary statistics of patients

Patient characteristics	Total	No PSE	PSE†	p Value*
Count of patient safety events	48 418	29 249	19 169	—
		60%	40%	—
Patient characteristics				
Average total charges (dollars)	18 493.36	11 268.06	29 518.90	0.000
SE	153.13	74.18	355.39	
Average length of stay (days)	5.59	3.71	8.46	0.000
SE	0.04	0.27	0.10	
Mean age	42.61	40.12	46.40	0.000
SE	0.11	0.14	0.17	
Complexity (scale=1–4) (%)				
0 (levels 1 and 2)	60.88	73.24	42.03	0.000
1 (levels 3 and 4)	39.12	26.76	57.97	
Sex (%)				
Male	49.23	49.17	49.32	0.685
Female	50.76	50.82	50.68	
Race (%)				
Caucasian	54.54	50.44	60.81	0.000
African—American	40.04	44.00	33.99	
Other	5.41	5.56	5.2	

*p Values calculated using t tests for continuous variables or χ^2 test for categorical variables.

†Includes non-preventable events such as adverse drug events and complications of care. PSE total contains duplicate events. PSE, patient safety event.

Table 4 Summary of case-finding results

Case-finding tool	Number of patient safety events (PSE) detected	PSE detected as percentage of total discharges (%)
E-codes	13 461	28
N-codes	13 253	27
Patient Safety Indicators	1463	3
Patient Safety Net	3596	7

In total, PSI detected 328 potential *puncture/laceration* events, N-codes detected 305, E-codes detected 192 and PSN detected two (figure 2). When linking *puncture/laceration* on unique identifiers, out of 447 potential events, 118 were identical between PSI and E-codes, 152 were identical between N-codes and E-codes and 188 were identical between PSI and N-codes. Only 100 events that were identified by PSI, E-codes and N-codes were identical. The two potential events detected by PSN were not among these 100.

DISCUSSION

This study investigated four tools for detecting potential patient safety events, identified the overlap among them and ascertained that multiple data sources and case-finding tools were necessary to maximise potential patient safety event surveillance. The example of *puncture/laceration* is illuminating. Out of a total of 447 potential events, there was an overlap of just 22% identified by three of the four methods (PSI, E-codes and N-codes), with two entirely distinct potential events identified by the fourth method, PSN.

Although it was not possible to conduct chart review, validation studies of the various case-finding tools give us confidence that the strong majority of the potential *puncture/laceration* events were appropriately identified. Gallagher's validation study of the *puncture/laceration* PSI conducted in 2005 found that 88.5% of cases identified were appropriate for quality-of-care investigation.¹¹ Similarly, Houchens *et al* found that the *puncture/laceration* PSI contained only 13% false positives that




Figure 2 Overlap of potential cases of puncture or laceration identified by four event detection tools. E-codes, external cause of injury; N-codes, nature of injury; PSI, Patient Safety Indicators; PSN, Patient Safety Net.

were POA, establishing this PSI's positive predictive value at 87%.⁸ In a 2009 report, Romano described even stronger performance by the puncture/laceration PSI, with 91% positive predictive value, and only 2% of false positives due to the injury being POA.²² The large percentage of hospital discharges flagged as potential patient safety events by E- and N-codes reflects the fact that these codes identify medical injuries reasonably well, but are less effective at distinguishing whether or not an injury is associated with medical error. A 1997 study in Victoria, Australia, found that E-codes could detect 65% of all adverse events identified by the chart review method.⁷ Similarly, a 2005 study in Wisconsin found that screening criteria based on E- and N-codes demonstrated 59.9% sensitivity and 97.4% specificity compared with medical record review.⁴ E- and N-code ranges were assigned to the most appropriate domain of our surveillance model, but many of these codes may or may not signal a medical error in any given case (figure 1). More definitive attributions of error must ultimately be made through investigation of individual cases, or through creation of new code ranges in the ICD-9-CM for medical misadventures.

This study is one of a small number to combine and compare the results of four or more detection tools applied to multiple linked datasets, and our findings are consistent with recent research showing the need to use more than one approach in patient safety surveillance.^{4 18 23}

The small amount of overlap among the four tools reflects the fact that each has different strengths and targets different occurrences. No case-finding tool is 'better' than another. Rather, one must understand the specific nature and utility of each tool. Owing to the variability in sensitivity and specificity of E- and N-coding, neither administrative data nor PSI should be used to compare the performance of different hospitals, though these data are extremely useful for internal quality assurance and improvement. By contrast, public health surveillance does not involve efforts to assign blame, or otherwise make individual institutions accountable for apparent gaps in patient safety. Public health surveillance is conducted from a population-based perspective and therefore does not require the level of accuracy that might be desired by clinicians. As Halperin and Baker point out, 'a surveillance system that does not have high sensitivity can still be useful in monitoring trends, as long as the sensitivity remains reasonably constant.'²⁴

Our results highlight the importance of including unique identifiers in surveillance data. Without them, discovering the degree of overlap would have been impossible. This is important because policy discussions of mandatory versus voluntary reporting of patient safety events have largely decided on submission of de-identified data, addressing legitimate privacy concerns. The absence of identifiers may limit the precision of national surveillance estimates.

Our study might have been strengthened by using chart review to verify the sensitivity and specificity of the four case-finding tools, but limited resources made such review prohibitive. Further validation of each tool is an important focus for future research. In addition, we applied our triangulation method to only one hospital. Future studies should consider this method with a larger sample of hospitals. Finally, advanced statistical methods to generate national patient safety event estimates must be developed.

It is well known that E-coding is not universal and its specificity is variable in the USA, limiting the generalisability of our findings.²⁵ There would be substantial public health benefit to regulatory change requiring uniform E-coding nationwide.

What is already known on this subject

- ▶ Error-associated medical injuries are estimated to cause 44 000–98 000 deaths per year in the USA.
- ▶ There is no existing surveillance system for medical injuries.
- ▶ Various case-finding tools exist, each providing insight into different aspects of the patient safety problem, but no coordination exists among them, and the resulting data are not integrated.

What this study adds

- ▶ This study combines and compares the results of multiple existing patient safety event detection tools applied to multiple linked datasets. Only a small amount of overlap was found, demonstrating the need for triangulation of data from different case-finding tools.
- ▶ A surveillance approach offers a more comprehensive design for the detection of patient safety events for public health assessment, policy development and evaluation.
- ▶ For research purposes, our results highlight the importance of including unique identifiers in surveillance data.

CONCLUSION

This study applies four different patient safety case-finding tools to multiple linked datasets. Even within the single domain of *error-associated injuries* analysed in this study, triangulation of multiple tools was needed to capture potential patient safety events more completely and, by eliminating duplication, more accurately. The case-finding tools used in this study are, to varying degrees, readily available: administrative data are standardised internationally, the PSI software for the SAS and Windows environment is available free of charge through the AHRQ website²⁶ and may be applied to any administrative dataset, and voluntary error-reporting systems, although they require some investment, can be set up by any hospital, healthcare network or national government. Our triangulation methodology may be applied in any country that uses these case-finding tools.

Triangulation through data linkage is a realistic trade-off between an ideal involving an unsustainable burden (chart review), and the acceptance of imperfect, but readily available data. Patient safety surveillance systems should be housed in a governmental institution providing affordable, easy access to the data, the analysis of which should produce new interventions. Comprehensive surveillance should provide the basis for optimal policy development and research funding, interventions and benchmarking to ensure we are making progress in improving patient safety.

Funding This research was supported (in part) by funding from the NIOSH Education and Research Center for Occupational Safety and Health at the Johns Hopkins Bloomberg School of Public Health—a doctoral training program (#T420H00842428).

Competing interests None.

Ethics approval This study was conducted with the approval of the Johns Hopkins Bloomberg School of Public Health IRB and Johns Hopkins Hospital.

Contributors All authors contributed to the conceptualisation, design and authorship of the study.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. **Kohn L**, Corrigan J, Donaldson M. *To Err is Human: Building a Safer Health System*. Washington, DC: National Academy Press, 2000.
2. **Dunn JD**. Error in medicine. *Ann Intern Med* 2001;**134**:342.
3. **Classen DC**, Lloyd RC, Provost L, *et al*. Development and evaluation of the institute for healthcare improvement global trigger tool. *J Patient Saf* 2008;**4**:169.
4. **Layde PM**, Meurer LN, Guse C, *et al*. Medical injury identification using hospital discharge data In *Advances in Patient Safety: From Research to Implementation*. Volume 2, AHRQ Publication No. 050021 (2). February 2005. Agency for Healthcare and Research Quality, Rockville, MD. <http://www.ahrq.gov/qual/advances/>.
5. **Utah Health Data Committee**. Adverse Events Related to Medical Care, Utah 1995–99. Salt Lake City, UT: Department of Health, 2001.
6. **Van Tuinen M**, Elder S, Link C, *et al*. Surveillance of Surgery-related Adverse Events in Missouri Using ICD-9-CM Codes. In *Advances in Patient Safety: From Research to Implementation*. Volume 1, AHRQ Publication No. 050021 (1). February 2005. Agency for Healthcare Research and Quality, Rockville, MD. <http://www.ahrq.gov/qual/advances/>.
7. **O'Hara DA**, Carson NJ. Reporting of adverse events in hospitals in Victoria, 1994–1995. *Med J Aust* 1997;**166**:460–3.
8. **Houchens RL**, Elixhauser A, Romano PS. How often are potential patient safety events present on admission? *Jt Comm J Qual Patient Saf* 2008;**34**:154–63.
9. **Iezzoni LI**. Finally present on admission but needs attention. *Med care* 2007;**45**:280.
10. **Agency for Healthcare Research and Quality**. AHRQ Quality Indicators - Guide to The Patient Safety Indicators. Agency for Healthcare Research and Quality, 2005. http://www.qualityindicators.ahrq.gov/psi_download.htm.
11. **Gallagher B**, Cen L, Hannan E. Validation of AHRQ's patient safety indicator for accidental puncture or laceration. *Adv Patient Saf* 2005;**2**:27–38.
12. **Bahl V**, Thompson MA, Kau TY, *et al*. Do the AHRQ Patient Safety Indicators flag conditions that are present at the time of hospital admission? *Med care* 2008;**46**:516.
13. **Glance LG**, Li Y, Osler TM, *et al*. Impact of date stamping on patient safety measurement in patients undergoing CABG: experience with the AHRQ Patient Safety Indicators. *BMC Health Serv Res* 2008;**8**:176.
14. **Romano PS**, Mull HJ, Rivard PE, *et al*. Validity of selected AHRQ patient safety indicators based on VA National Surgical Quality Improvement Program data. *Health Serv Res* 2009;**44**:182–204.
15. **Murff HJ**, Patel VL, Hripsak G, *et al*. Detecting adverse events for patient safety research: a review of current methodologies. *J Biomed Inform* 2003;**36**:131–43.
16. **Berenholtz SM**, Pronovost PJ. Monitoring patient safety. *Crit Care Clin* 2007;**23**:659–73.
17. **Agency for Healthcare Research and Quality**. *Reducing and Preventing Adverse Drug Events To Decrease Hospital Costs*. Research in Action 2001. <http://www.ahrq.gov/qual/aderia.htm> (accessed 17 Nov 2010).
18. **Naessens JM**, Campbell CR, Huddlestone JM, *et al*. A comparison of hospital adverse events identified by three widely used detection methods. *Int J Qual Health Care* 2009;**21**:301–7.
19. **Miller MR**, Elixhauser A, Zhan C, *et al*. Patient Safety Indicators: using administrative data to identify potential patient safety concerns. *Health Serv Res* 2001;**36**:110–32.
20. **Zhan C**, Miller MR. Excess length of stay, charges, and mortality attributable to medical injuries during hospitalization. *JAMA* 2003;**290**:1868–74.
21. **Romano PS**, Geppert JJ, Davies S, *et al*. A national profile of patient safety in U.S. hospitals. *Health Aff (Millwood)* 2003;**22**:154–66.
22. **Romano PS**. *Lessons from the AHRQ PSI Validation Pilot Project*. Slide Presentation from the AHRQ 2008 Annual Conference (Text Version). Agency for Healthcare Research and Quality, 2009. <http://www.ahrq.gov/about/annualmtg08/091008slides/Romano.htm>.
23. **Samore MH**, Evans RS, Lassen A, *et al*. Surveillance of medical device-related hazards and adverse events in hospitalized patients. *JAMA* 2004;**291**:325–34.
24. **Halperin W**, Baker E. *Public Health Surveillance*. New York: Van Nostrand Reinhold, 1992.
25. **Annest JL**, Fingerhut LA, Gallagher SS, *et al*. Strategies to improve external cause-of-injury coding in state-based hospital discharge and emergency department data systems: recommendations of the CDC Workgroup for Improvement of External Cause-of-Injury Coding. *MMWR Recomm Rep* 2008;**57**:1–15.
26. **Inoue K**, Hirotsawa I, Yatsuduka M, *et al*. Utilization of a voluntary reporting system in quantitative risk assessment for medical tasks in a hospital setting—with special reference to tasks done by nurses. *J Occup Health* 2002;**44**:360–72.