Conclusion Specific road traffic environments raise the severity of distracted walking, which further threatens pedestrian safety.

Learning Outcomes Specific road environmental risk increases the severity of distracted walking, additionally creating an indirect impact on near-crash event.

2D.004 PEDESTRIANS' STREET CROSSING PERCEPTIONS ON PEDESTRIAN BRIDGES IN DAR ES SALAAM

^{1,2,3}Daudi Katopola*, ²Fredrick Mashili, ¹Marie Hasselberg. ¹Department of Global Public Health, Stockholm, Sweden; ²Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania; ³National Institute of Transport, Dar es Salaam, Tanzania

10.1136/injuryprev-2021-safety.47

Background About 30 per cent of all road traffic incidence deaths in Tanzania involve pedestrians. Previous studies show that pedestrian bridges might not necessarily result in high bridge use, even when pedestrians must cross 3–4 lanes with heavy traffic. This paper explores the perceptions of pedestrians when crossing urban roads through using or not using pedestrian bridges.

Methods Nineteen semi-structured interviews were purposively conducted around six pedestrian bridges within Dar es Salaam. All interviews were conducted in Swahili, recorded using digital devices, transcribed verbatim then translated into English. To know the perceptions of pedestrians with respect to their experiences of using or not using the bridges, content analysis was employed with the help of a qualitative data analysis software (MAXQDA).

Results Three key themes were identified, relating to planning and decision making, alternative activities carried out around pedestrian bridges, and behavioural intentions and perceptions of pedestrians when accessing the bridges. Results also suggest that, participants are aware that the aim of the bridges are to prevent road traffic injuries. Conversely, participants were sceptical about inappropriate behavioural use of the bridges when linked to planning decisions and norms.

Conclusion The findings support the idea that already vulnerable road users such as pregnant women, older people and physically challenged pedestrians are exposed to higher risks of being injured since they cannot use the bridges when crossing roads.

Learning Outcomes Involvement of users is crucial when constructing pedestrian bridges in order to increase awareness on appropriate behavioural use when crossing roads with bridges.

2D.005 OLDER PEDESTRIAN INJURY OUTCOMES: A SYSTEMATIC REVIEW, META-ANALYSIS, AND GRADE ASSESSMENT

JE Rod*, Oscar Oviedo-Trespacios, Teresa Senserrick, Mark King. *CARRS-Q, QUT, Brisbane, Australia*

10.1136/injuryprev-2021-safety.48

Background Older pedestrians 60+ (OP) have worse health outcomes due to pedestrian trauma. There is no available systematic evaluation of OP injury outcomes. This becomes increasingly important as the global population ages. We conducted a systematic review, meta-analysis and GRADE

assessment of OP injury severity and compare them with those of younger age groups.

Methods We searched PubMed, Embase, Scopus, CINAHL, PsycInfo, AMED, Web of Science, LILACS and TRID to identify studies assessing OP crashes and falls. Reporting was evaluated using STROBE, and GRADE was used to assess evidence quality. Random-effect model meta-analysis served to obtain pooled estimates. The review is registered in PROSPERO (#CRD42019140786).

Results We included 60 studies (1,012,041 pedestrians). The pooled incidence proportion of OP crashes was 25% (95% CI: 22.3 – 28.5). OP were found to have higher severe injury, more critical care admissions, and higher fatality than younger pedestrians. OP also have higher incidence rates and severe injury from pedestrian falls. The overall quality of the evidence supporting the findings was low.

Conclusions OP risks may offset some of the health gains from active ageing strategies, especially for developing economies with high numbers of vulnerable road users and with rapid population ageing.

Learning Outcomes There is a need for research to improve the quality of evidence in regard to short and medium-term health outcomes and address long-term health outcomes for OP trauma.

2E – Intentional, March 23, 2021

2E.001 MEASURING THE EFFECTIVENESS OF PREVENTION OF FAMILY VIOLENCE IN A REGIONAL SETTING

¹Margie Robinson*, ¹Heath Greville, ¹John Woods, ²Barbara Nattabi, ¹Monica Moran, ²Colleen Fisher, ¹Sandra Thompson, ²Karen Martin. ¹WA Centre for Rural Health, School of Population and Global Health, University of Western Australia, Geraldton, Australia; ²School of Population and Global Health, University of Western Australia, Nedlands, Australia

10.1136/injuryprev-2021-safety.49

Background Rates of family and gender-based violence (FV) are high, particularly in rural and remote areas in Australia. The City of Greater Geraldton, a local government area in the Midwest region of Western Australia, has developed a FV primary prevention strategic action plan. Local data are needed to measure the impact of community based prevention efforts. Our research aim was to develop a tool to measure community attitudes and experiences of FV in order to assess the effectiveness of efforts to change attitudes and reduce violence over time.

Methods A working group developed the Local Community Attitudes and Exposure to Violence Survey (LCAEVS), based primarily upon the National Community Attitudes Survey (NCAS), which is periodically undertaken by Australia's National Research Organisation for Women's Safety (ANROWS). Questions from all relevant domains in the NCAS were included, with prioritisation of items likely to be sensitive to change over time. The survey was shortened to reduce respondent burden.

Results Geraldton respondents showed misunderstandings about the key drivers and the gendered pattern of FV, while some local attitudes were less supportive of gender equality than those documented in the national survey.

Conclusions Approaches adopted to distribute the survey, the response rate and demographics of the population who completed it, and comparison of responses of Geraldton residents with those in the NCAS will be discussed.

Learning Outcomes It is feasible to administer a shorter survey based on the NCAS, to measure particular attitudes on which to focus violence prevention education for a regional population.

2E.002 CYCLE OF VIOLENCE: CHILDHOOD ABUSE AND RISK OF VIOLENCE REVICTIMISATION IN ADULTHOOD

¹Mark A Bellis, ²Nadia Butler*, ²Zara Quigg. ¹Public Health Wales, Wrexham, UK; ²Liverpool John Moores University, Liverpool, UK

10.1136/injuryprev-2021-safety.50

Interpersonal violence is a serious threat to the attainment of the Sustainable Development Goals (SDGs). A public health approach to violence prevention is crucial, and addressing risk factors is a key priority. Global research has demonstrated that childhood adversity increases risk of a range of poor outcomes across the lifecourse. This study used data from a nationally representative survey of household residents (n=21,845), to examine the impact of childhood abuse (physical, sexual and psychological abuse, and witnessing domestic violence) on risk of adulthood violence revictimisation (physical assault (PA), intimate partner violence (IPV), and sexual violence (SV)).

Compared to individuals who experienced no child abuse, those who experienced one type were, twice as likely to experience PA, and three times as likely to have experienced IPV and/or SV. Individuals who experienced multiple types were three, six and seven times more likely to experience PA, IPV, and SV, respectively. After controlling for the number of types experienced, associated types differed by adult violence outcome; child psychological and physical abuse were associated with IPV; psychological and sexual abuse with SV; and psychological abuse with PA.

Findings from the study will be presented with consideration of strategies to prevent and respond to child abuse and the potential downstream effect on preventing interpersonal violence across the lifecourse and achieving the SDGs. With adulthood victimisation likely to compound the already detrimental effects of childhood abuse, and given that many associated outcomes represent adversities for the next generation, breaking the cycle of violence represents a critical priority.

2E.003 BURDEN OF SUICIDE IN NEPAL: AN ANALYSIS OF POLICE RECORDS

¹Bhagabati Sedain Silwal*, ²Puspa Raj Pant., ³Yogendra Bahadur Gurung. ¹Tribhuvan University Padmakanya Multiple Campus, Kathmandu, Nepal; ²1.Nepal Injury Research Centre (NIRC) 2 University of the West of England (UWE Bristol), Bristol, UK; ³Tribhuvan University Central Department of Population Studies, Kathmandu, Nepal

10.1136/injuryprev-2021-safety.51

Background Suicide is a complex human behaviour which might be initiated with an onset of psychic pain. It is the leading cause of unnatural deaths in Nepal, and the number is rising continuously. Due to issues regarding the legality, social

stigma, and logistical problems, there is an underreporting of suicide cases in Nepal.

Methods We studied the records of Nepal Police for five years between 2015 and 2019. Access to the dataset was obtained through institutional research collaborations. The socio-demographic variables for each individual case of deaths were extracted and analysed.

Results During the five years, 25,316 cases of suicidal deaths were recorded; an average of 14 people each day. The data revealed that people aged 19 to 35 years accounted for almost 60% of all suicide deaths. Overall, there is an increase in suicidal deaths by 33% in 2019 compared to 2015 whereas the increase in females was 200%. Hanging was the most common (70.13%) method followed by the consumption of poison (26.55%).

Conclusion Suicide is a serious but neglected public health problem in Nepal. It has been found that adolescents, youths, and females are the most vulnerable population for suicide. The findings warrant an urgent need for suicide prevention in Nepal.

Learning Outcomes Police records are currently the only reliable source for information regarding suicides occurring within the nation. There still may be cases that might not have been recorded. Large-Scale researches can identify factors associated with suicides in Nepal in order to inform prevention interventions.

2E.004 ZERO SUICIDE HEALTHCARE: PROGRAM THEORY TO GUIDE EVALUATION FOR SAFER SUICIDE CARE

¹Sue Murray*, ²Alan Woodward*. ¹Zero Suicide Institute of Australasia, Balmain, Australia; ²Alan Woodward Consulting, Kiama, Australia

10.1136/injuryprev-2021-safety.52

Context The World Health Organisation estimates one person dies by suicide every 40 seconds. In Australia it's estimated around 20% of suicides are people who have been in the Australian healthcare system. This is not an acceptable outcome for modern healthcare systems.

Process The Zero Suicide Healthcare (ZSH) framework has been adopted in many developed countries, comprised of seven elements designed to build organisational capability, improve clinical practice and create better service pathways for recovery.

Analysis ZSH draws on techniques of quality management and continuous improvement. It implicitly assumes suicide prevention can be addressed in health care settings in the same way, and with the same absolute improvements, as has been done in wound management, infection control and medication management. It advances evidence-based practice in suicide prevention as a standardised and systematic reform. It encourages the use of data and measurement to monitor performance for continuous improvement within healthcare settings.

However, there is no single source of understanding of how the elements of the ZSH framework interact to achieve outcomes. The development of the Theory of Change for ZSH provides greater clarity about the overall design of the framework using program theory. This will also underpin the evaluation of ZSH across projects.

Outcomes A model will be presented demonstrating alignment to the ZSH framework and giving a consistent approach to