laboratory workers to hidden and unnecessary hazards. Even having the prospect of working in a new purpose-built facility does not mean that the laboratory worker is not a risk from the hidden hazards which form part of the architects/designer need to design an aesthetically pleasing workspace.

Perhaps we could forgive the repurposed laboratory, given limitations in budget, and infrastructure, but there are no excuses when it comes to the design and development of new facilities. Whether it is a new 'super laboratory' or a small QA/Formulation laboratory, it is important that designers/architects understand some of the basics concepts which are essential for developing a safe workplace. This includes understanding the role that AS/NZS 2243 Safety in Laboratories and AS/NZS 2982 Laboratory design and construction have in designing laboratory and auxiliary workspaces (Store rooms, preparation rooms).

E-Posters P2 – Intentional, Trauma, Violence, Suicide, Poisoning & Alcohol, March 23, 2021

P2.001 ALCOHOL'S EXCESSIVE EPISODIC CONSUMPTION IN TEENAGERS FROM EL CARMEN, JUJUY, ARGENTINA, 2018

Nadia Rivera Leinecker*. Faculty of humanities and social sciences. Jujuy national university, El Carmen, Argentina

10.1136/injuryprev-2021-safety.91

The youth population of Jujuy has adopted a pattern of consumption with large amounts of alcohol in short periods of time, called excessive episodic consumption of alcohol or 'binge drinking', which is intake of large amounts of alcohol (more than 5 UNEs = 60 Gr/cc pure alcohol) in a short period of time (hours). This produces significant behavior and consciousness state alterations and causes negative social-sanitary effects.

Design Quantitative, transversal descriptive research. Applying structured, self-administered, anonymous and voluntary questionnaire.

Population and sample Intentional sampling of students in the first and last high school's years of educational centers in El Carmen.

Results Excessive episodic consumption was more prevalent among older students (28.0% vs. 9.2%); among those who worked (26.0% vs. 15.8%); who had friends who drank alcohol (29.1% vs. 5.8%); that were concerned about their consumption (48.3% vs. 15.8%); who said they had sufficient information on alcohol use (21.0% vs. 12.6%) or on alcohol abuse (21.9% vs. 11.6%); who regularly attended bars or pubs (36.6 vs. 12.9%); and among those who spent a lot of time with their partner (22.8% vs. 13.4%). Excessive episodic consumption was lower among those who spent a lot of time watching television (15.2% vs. 25.6%); with their family (14.8% vs. 23.7%); who spent time in the church (14.1% vs. 20.7%); or in some shops where playing video games (12.9% vs. 19.0%).

Conclusion Excessive alcohol consumption is enhanced by having friends who drink and attending night bars.

Reflections Strategies for risk reduction in consumption contexts are necessary.

P2.002 TODDLER INTERACTION WITH LIQUID LAUNDRY DETERGENT CAPSULES: INFLUENCE OF CAPSULE SIZE

¹Annalise Richmond*, ²David Schwebel, ²Casey Morgan, ¹Gerard Stijntjes. ¹Procter and Gamble, Research and Development, Srombeek Bever, Belgium; ²Department of Psychology, University of Alabama at Birmingham, Birmingham, USA

10.1136/injuryprev-2021-safety.92

Background Liquid laundry capsules have been involved in child poisoning incidents, allegedly due to product appearances. This research explored potential influence of capsule size on toddler interaction.

Methods We compared two capsule designs identical in appearance but different in size through three studies: (a) forced-choice test in laboratory setting; (b) simulated real-world setting of laundry cabinet with open container of capsules; and (c) simulated real-world cabinet with capsule outside its container. Capsule interaction was measured by grasping choice among samples of 156 toddlers ages 9–36 months in each study.

Results Study 1: Toddlers selected the small (49.8% selection) and large (50.2%) capsule with nearly identical frequency. Study 2: Toddlers grasped the small (26.8%) and large (22.3%) capsule or container of capsules with similar frequency. Study 3: Toddlers grasped the small (18.0%) and large (19.2%) capsules with similar frequency. No consistent gender or age differences existed. Study 1 found toddlers selected capsules closest to their dominant hand more frequently, no matter their size.

Conclusion Findings consistently failed to demonstrate statistical differences in toddler interaction with different-sized capsules. We conclude the difference in size (small vs large capsules) is not expected to impact toddler interaction rates in the home.

Learning Outcomes Extending previous research concerning laundry capsule colours, size also seems unlikely to impact child poisoning risk. The work offers valuable evidence for action to reduce child poisoning incidents, supporting current strategies to reduce risk through theory-driven caregiver education and use of packaging with barriers to prevent young children's access to capsules.

P2.003 CHARACTERISTICS OF INTERNET-BASED MEDIA REPORTS ABOUT INFLUENTIAL VIOLENCE AGAINST DOCTORS IN CHINA

¹Liheng Tan, ¹Shujuan Yuan, ¹Peixia Cheng, ¹Peishan Ning, ¹Yuyan Gao, ¹Wangxin Xiao, ³David Schwebel, ¹Guoqing Hu*. ¹Department of Epidemiology and Health Statistics, Xiangya School of Public Health, Central South University, Changsha, China; ²Kaifu District Center for Disease Control and Prevention, Changsha, China; ³Department of Psychology, University of Alabama at Birmingham, Birmingham, USA

10.1136/injuryprev-2021-safety.93

Background Media reports may influence criminal behavior that leads to violence against doctors. Little is known about the characteristics of media reports on patient violence against their doctors.