

Table S3: Comparison of Pooled Cohort Study PedsQL Non-participants (Missing) and Participants (Present)*

	Missing	Present	Total N=2436
Sex			
Male	119 (67.2)	1493 (66.1)	1612 (66.2)
Female	58 (32.8)	766 (33.9)	824 (33.8)
Age group			
0-4 years	51 (28.8)	619 (27.4)	670 (27.5)
5-9 years	38 (21.5)	523 (23.2)	561 (23.0)
10-14 years	56 (31.6)	754 (33.4)	810 (33.3)
15-17 years	32 (18.1)	363 (16.1)	395 (16.2)
Socio Economic Status (SES)			
Low SES	81 (49.4)	823 (38.2)	904 (39.0)
Moderate SES	56 (34.1)	898 (41.7)	954 (41.1)
High SES	27 (16.5)	434 (20.1)	461 (19.9)
Not available	13	104	117
Country			
VSTR	56 (31.6)	910 (40.3)	966 (39.7)
CHAI	35 (19.8)	889 (39.4)	924 (37.9)
UKBOI	16 (9.0)	165 (7.3)	181 (7.4)
BCCH-LIO	70 (39.5)	295 (13.1)	365 (15.0)
Transport status			
Non-transport	119 (77.8)	1620 (74.2)	1739 (74.5)
Transport	34 (22.2)	562 (25.8)	596 (25.5)
Not available	24	77	101
Hospital or ED			
ED only	73 (41.2)	880 (39.0)	953 (39.2)
Hospital admission	104 (58.8)	1377 (61.0)	1481 (60.8)
Not available	-	<5	<5
Comorbidity status			
No comorbidities	158 (89.3)	2033 (90.0)	2191 (89.9)
At least 1 comorbidity	19 (10.7)	226 (10.0)	245 (10.1)
Injury Severity Score (ISS) Tertiles			
Low (1-4)	86 (50.9)	957 (43.8)	1043 (44.3)
Mid (5-16)	38 (22.5)	591 (27.1)	629 (26.7)
High (17+)	45 (26.6)	635 (29.1)	680 (28.9)
Not available	8	76	84
Intent			
Unintentional	149 (84.2%)	2054 (90.9%)	2203 (90.4%)
Intentional	13 (7.3%)	108 (4.8%)	121 (5.0%)
Intent not known	15 (8.5%)	97 (4.3%)	112 (4.6%)
Injury (any)			
N19, N26 Fracture of femur	11 (6.2)	107 (4.7)	118 (4.8)
N20 Fracture of patella, tibia, fibula, or ankle	6 (3.4)	102 (4.5)	108 (4.4)
N28 Moderate to severe traumatic brain injury	41 (23.2)	611 (27.0)	652 (26.8)
N37, N17, N18 Crush injury, fracture foot/hand bones	6 (3.4)	78 (3.5)	84 (3.4)
N43 Internal hemorrhage in abdomen or pelvis	18 (10.2)	265 (11.7)	283 (11.6)
N27 Minor traumatic brain injury	16 (9.0)	626 (27.7)	642 (26.4)
N21 Fracture of pelvis	7 (4.0)	102 (4.5)	109 (4.5)

N42 Severe chest Injury	16 (9.0)	193 (8.5)	209 (8.6)
N8, N9, N10 Burns (including lower airways)	11 (6.2)	52 (2.3)	63 (2.6)
N25 Fracture of vertebral column	5 (2.8)	104 (4.6)	109 (4.5)
N35, N36 Asphyxiation, Non-fatal submersion	<5	<20	<20
N40, N44 Contusion, open wound	31 (17.5)	432 (19.1)	463 (19.0)
N14 Other injuries of muscle & tendon and other dislocations	27 (15.3)	190 (8.4)	217 (8.9)
N15 Fracture of clavicle, scapula, or humerus	9 (5.1)	218 (9.7)	227 (9.3)
N22 Fracture of radius or ulna	16 (9.0)	238 (10.5)	254 (10.4)
Other	68 (38.4)	1003 (44.4)	1071 (44.0)

Total missing = 7%. *Small bases reduced to ensure privacy.