

**Table S3:** Comparison of Pooled Cohort Study PedsQL Non-participants (Missing) and Participants (Present)\*

	Missing	Present	Total N=2436
<b>Sex</b>			
Male	119 (67.2)	1493 (66.1)	1612 (66.2)
Female	58 (32.8)	766 (33.9)	824 (33.8)
<b>Age group</b>			
0-4 years	51 (28.8)	619 (27.4)	670 (27.5)
5-9 years	38 (21.5)	523 (23.2)	561 (23.0)
10-14 years	56 (31.6)	754 (33.4)	810 (33.3)
15-17 years	32 (18.1)	363 (16.1)	395 (16.2)
<b>Socio Economic Status (SES)</b>			
Low SES	81 (49.4)	823 (38.2)	904 (39.0)
Moderate SES	56 (34.1)	898 (41.7)	954 (41.1)
High SES	27 (16.5)	434 (20.1)	461 (19.9)
Not available	13	104	117
<b>Country</b>			
VSTR	56 (31.6)	910 (40.3)	966 (39.7)
CHAI	35 (19.8)	889 (39.4)	924 (37.9)
UKBOI	16 (9.0)	165 (7.3)	181 (7.4)
BCCH-LIO	70 (39.5)	295 (13.1)	365 (15.0)
<b>Transport status</b>			
Non-transport	119 (77.8)	1620 (74.2)	1739 (74.5)
Transport	34 (22.2)	562 (25.8)	596 (25.5)
Not available	24	77	101
<b>Hospital or ED</b>			
ED only	73 (41.2)	880 (39.0)	953 (39.2)
Hospital admission	104 (58.8)	1377 (61.0)	1481 (60.8)
Not available	-	<5	<5
<b>Comorbidity status</b>			
No comorbidities	158 (89.3)	2033 (90.0)	2191 (89.9)
At least 1 comorbidity	19 (10.7)	226 (10.0)	245 (10.1)
<b>Injury Severity Score (ISS) Tertiles</b>			
Low (1-4)	86 (50.9)	957 (43.8)	1043 (44.3)
Mid (5-16)	38 (22.5)	591 (27.1)	629 (26.7)
High (17+)	45 (26.6)	635 (29.1)	680 (28.9)
Not available	8	76	84
<b>Intent</b>			
Unintentional	149 (84.2%)	2054 (90.9%)	2203 (90.4%)
Intentional	13 (7.3%)	108 (4.8%)	121 (5.0%)
Intent not known	15 (8.5%)	97 (4.3%)	112 (4.6%)
<b>Injury (any)</b>			
N19, N26 Fracture of femur	11 (6.2)	107 (4.7)	118 (4.8)
N20 Fracture of patella, tibia, fibula, or ankle	6 (3.4)	102 (4.5)	108 (4.4)
N28 Moderate to severe traumatic brain injury	41 (23.2)	611 (27.0)	652 (26.8)
N37, N17, N18 Crush injury, fracture foot/hand bones	6 (3.4)	78 (3.5)	84 (3.4)
N43 Internal hemorrhage in abdomen or pelvis	18 (10.2)	265 (11.7)	283 (11.6)
N27 Minor traumatic brain injury	16 (9.0)	626 (27.7)	642 (26.4)
N21 Fracture of pelvis	7 (4.0)	102 (4.5)	109 (4.5)

---

N42 Severe chest Injury	16 (9.0)	193 (8.5)	209 (8.6)
N8, N9, N10 Burns (including lower airways)	11 (6.2)	52 (2.3)	63 (2.6)
N25 Fracture of vertebral column	5 (2.8)	104 (4.6)	109 (4.5)
N35, N36 Asphyxiation, Non-fatal submersion	<5	<20	<20
N40, N44 Contusion, open wound	31 (17.5)	432 (19.1)	463 (19.0)
N14 Other injuries of muscle & tendon and other dislocations	27 (15.3)	190 (8.4)	217 (8.9)
N15 Fracture of clavicle, scapula, or humerus	9 (5.1)	218 (9.7)	227 (9.3)
N22 Fracture of radius or ulna	16 (9.0)	238 (10.5)	254 (10.4)
Other	68 (38.4)	1003 (44.4)	1071 (44.0)

---

Total missing = 7%. \*Small bases reduced to ensure privacy.